Synthesis of Radioiodinated Carbocyclic Cytosine Analogues

Hyunseok Ahn,[†] Gwangil An,[‡] and Hakjune Rhee^{†,§,*}

[†]Department of Bionanotechnology, Hanyang University, Kyunggi-do 426-791 Korea. *E-mail: hrhee@hanyang.ac.kr

[‡]Laboratory of Radiopharmaceuticals and Laboratory of Nuclear Medicine, Seoul 139-706, Korea

[§]Department of Chemistry and Applied Chemistry, Hanyang University, Kyunggi-do 426-791 Korea

Received March 29, 2011, Accepted April 21, 2011

The synthesis of carbocyclic analogues of normal nucleosides has grown exclusively since they have shown potential antiviral and antitumor activities. Radiolabeled *cis*-1-[4-(hydroxy-methyl)-cyclopent-2-enyl]-5-^[124I]-iodocytosine (carbocyclic d4IC) and *cis*-1-[4-(hydroxy-methyl)-cyclopent-2-enyl]-5-(2-^[124I]iodovinyl)cytosine (carbocyclic d4IVC) were synthesized. The synthetic route employed Pd(0)-catalyzed coupling reaction together with organotin and exchange reaction for radioiodination as key reactions. Carbocyclic ^[124I]d4IC gave more than 75% radiochemical yield with greater than 95% radiochemical purity. Carbocyclic ^[124I]d4IVC gave more than 80% radiochemical yield with greater than 95% radiochemical purity.

Key Words : Carbocyclic nucleoside, Radioiodine, PET, Radiopharmaceutical, Labeling

Introduction

Herpes Simplex Virus type-1 thymidine kinase (HSV1-tk) is widely studied for the imaging of gene expression.¹ Various radiolabeled nucleoside analogues, both pyrimidine and acycloguanosine derivatives, have been synthesized and evaluated for HSV1-tk imaging.² Normal nucleoside analogues have lesser metabolic stability. The glycosidic linkage in normal nucleosides is cleaved by thymidine phosphorylase enzymes.³ The replacement of furanose oxygen by carbon gives the nucleoside greater stability *in vivo*. Because of this metabolic stability, carbocyclic nucleosides and their analogues have been investigated by many research groups.^{4,5} Recently, our group reported synthesis and biological evaluation of uracil based radioiodinated carbocyclic nucleosides, *cis*-1-[4-(hydroxymethyl)-cyclopent-2-enyl]-5-^[124I]iodouracil (carbocyclic d4IU, **1**) and *cis*-1-[4-(hydroxymethyl)-cyclopent-2-enyl]-5-(2-^[124I]iodovinyl)uracil (carbocyclic d4IVU, **2**) (Figure 1).⁶ Since the previous result showed the radioiodine labeled carbocyclic d4IU (**1**) and d4IVU (**2**) are potent imaging materials for HSV1-tk, we decided to extend our research to synthesize cytosine-based

carbocyclic nucleosides. Here, we report the synthesis of cytosine-based radiolabeled carbocyclic nucleosides, *cis*-1-[4-(hydroxymethyl)-cyclopent-2-enyl]-5-^[124I]iodocytosine (carbocyclic d4IC, **3**) and *cis*-1-[4-(hydroxymethyl)-cyclopent-2-enyl]-5-(2-^[124I]iodovinyl)cytosine (carbocyclic d4IVC, **4**) as potential HSV1-tk imaging reporter probes (Figure 1).

Results and Discussion

In recent years, our group has synthesized carbocyclic nucleosides and their analogues.⁵⁻⁷ We have prepared dicarbonate **10** (Scheme 1) as a key intermediate. In this paper, we used this dicarbonate **10** again to synthesize new carbocyclic nucleosides. Synthesis of the dicarbonate **10** was carried out using known procedures with a minor modification. For the ring-opening step of hydroxylactone **7** to triol **8**, sodium borohydride was used.⁸ This method has an advantage in having a simpler and faster work-up than when using lithium aluminumhydride.⁵

Compounds **10** and 5-iodocytosine (**11**) were coupled by a

Scheme 1. (a) Toluene, H₂O, rt, 12 h, 40% (b) NaBH₄, CH₃OH, rt, 1 h, 90%. (c) (i) NaIO₄, diethyl ether, H₂O, rt, 2 h (ii) ethylene glycol, rt, 1 h (iii) NaBH₄, rt, 2 h, 70%. (d) methyl chloroformate, DMAP, CHCl₃, pyridine, 0 °C, 1 h, 93%.

Figure 1. Carbocyclic ^[124I]d4IU (**1**), ^[124I]d4IVU (**2**), ^[124I]d4IC (**3**) and ^[124I]d4IVC (**4**).

Scheme 2. (a) (i) $\text{Pd}(\text{OAc})_2$, $(i\text{-PrO})_3\text{P}$, THF, under argon, rt (ii) $n\text{-BuLi}$, hexane, rt. (iii) **11** and NaH in DMSO (iv) **10** in anhydrous THF, 12 h, 70%. (b) (i) $\text{Pd}_2(\text{dba})_3$, DMF (ii) bis(tributyltin), 100 °C, 7 h, 47%. (c) (i) 0.5 *N* aqueous K_2CO_3 , rt, 12 h (ii) 1.0 *N* HCl to pH 7.0, 71%.

$\text{Pd}(0)$ -catalyzed Tsuji-Trost cross-coupling reaction.⁹ The $\text{Pd}(0)$ catalyst was prepared *in situ* using $\text{Pd}(\text{OAc})_2$ in THF with triisopropyl phosphite and $n\text{-BuLi}$. The produced π -allylpalladium complex of compound **10** undergoes nucleophilic attack by the anion on the pyrimidine base **11**.⁵⁻⁷ Radiolabeling reactions with organotin compounds have been reported as a facile strategy.^{6,10} Thus, we applied this method for our synthesis. Compound **12** was converted to the desired stannylated precursor **13** by the $\text{Pd}(0)$ -catalyzed condensation of bis(tributyltin).⁶ In addition, compound **12** was hydrolyzed to produce *cis*-1-[4-(hydroxymethyl)-2-cyclopenten-1-yl]-5-iodocytosine (**14**) as a cold form of the reference compound.

For synthesis of the iodovinyl compound, a trialkylsilylvinyl precursor was selected.^{7,11} Compound **12** was converted to a trimethylsilylvinyl compound **15** by using a $\text{Pd}(0)$ -catalyzed Stille coupling reaction.^{7,12} After the hydrolysis of compound **15** and the following exchange reaction of compound **16** with ICl , compound **17** was obtained as a cold form of reference material.⁷

As shown in Scheme 4, the radioiodination of **13** was performed using a no-carrier-added iodine-124 produced by the Korea Institute of Radiological and Medical Sciences (KIRAMS). Labeled compound **18** was hydrolyzed by 0.5 *N*

Scheme 4. (a) (i) Na^{124}I , 0.01 *N* NaOH, MeOH (ii) 1.0 *N* HCl to pH 4.0, 30% H_2O_2 , 10 min (iii) NaHSO_3 . (b) (i) 1.0 *N* K_2CO_3 , rt, 1 h (ii) 1.0 *N* HCl to pH 7.

potassium carbonate. Then the desired carbocyclic d4IC (**3**) was separated, collected, and analyzed using reverse phase HPLC with UV and radioactivity detectors. The HPLC chromatographic retention time of carbocyclic d4IC (**3**) was 19.65 min with a radiochemical yield > 75% and radiochemical purity > 95%. We confirmed that no side products were produced except for the unreacted radioiodide based on radio-TLC monitoring. Moreover, radiolabeled and reference compounds have the same retention time in HPLC.

With the same method, compound **16** was radioiodinated, separated, collected, and analyzed using reverse phase HPLC with UV and radioactivity detectors. The HPLC chromatographic retention time of carbocyclic d4IVC (**4**) was 17.0 min with a radiochemical yield > 80% and radiochemical purity > 95%.

Scheme 3. (a) $(E)\text{-Bu}_3\text{SnCH=CHSiMe}_3$, $\text{Pd}_2(\text{dba})_3$, DMF, 12 h, 60 °C, 60%. (b) (i) 0.5 *N* aqueous K_2CO_3 , rt, 12 h (ii) 1.0 *N* HCl, 93%. (c) ICl , CH_3CN , 30 min, 70%.

Scheme 5. (a) (i) Na^{124}I , 0.01 *N* NaOH, MeOH (ii) 1.0 *N* HCl to pH 4.0, 30% H_2O_2 , 10 min (iii) NaHSO_3 .

Conclusions

We successfully synthesized radioiodinated *cis*-1-[4-(hydroxymethyl)-cyclopent-2-enyl]-5-[¹²⁴I]iodocytosine (carbocyclic d4IC, **3**) and *cis*-1-[4-(hydroxymethyl)-cyclopent-2-enyl]-5-(2-[¹²⁴I]iodovinyl)cytosine (carbocyclic d4IVC, **4**). The biological evaluation of these compounds will be reported in due course.

Experimental

General. ¹H and ¹³C NMR spectra were obtained using a Bruker 300 spectrometer (300 MHz) and a Varian 500 spectrometer (500 MHz). Infrared spectra were recorded on a Bruker Alpha FT-IR spectrometer. Uncorrected melting points were determined with a Sanyo Gallenkamp melting point apparatus. HRMS were obtained with a JMS 700 spectrometer. Analytical thin layer chromatography (TLC) was conducted on E. Merck 60 F254 aluminum-backed silica-gel plates (0.2 mm) with a fluorescent indicator. Radio-TLC was measured with a Bioscan AC-3000 scanner (Washington D.C.). High performance liquid chromatography (HPLC) was carried out on a μ Bondapak C18 column (7.8 $\times$ 300 mm, Waters). Developed plates were visualized with UV light or a 2.0% phosphomolybdic acid staining solution. Flash column chromatography was performed using Merck silica-gel 60 (230-400 mesh) under positive pressure. All reagents and solvents were reagent grade and purified by known procedures before use. For radioiodine labeling, no-carrier-added iodine-124 was produced *via* the ¹²⁵Te(p, 2n) ¹²⁴I nuclear reaction in an enriched ¹²⁵TeO₂ at the KIRAMS MC-50 cyclotron.

5-(1,2-Dihydroxyethyl)-2-cyclopentenol (8**).**⁵ Sodium borohydride (17.0 g, 450 mmol) was slowly added to a solution of **7** (21.0 g, 150 mmol) in anhydrous methanol (1.0 L) in an ice bath. After stirring the resulting solution for 1 h at ambient temperature, the solvent was removed by rotary-evaporation. The residue was diluted with water and neutralized with 1.0 *N* aqueous HCl to reach a pH of 7. The aqueous solution was concentrated by rotary-evaporation. The residue was diluted with ethyl acetate, dried with anhydrous MgSO₄, and concentrated by rotary-evaporation. The crude mixture was purified by silica-gel column chromatography (CH₂Cl₂:MeOH = 7:1, v/v) to give colorless oil **8** (*R*_f = 0.25; 15.1 g, 70%). ¹H NMR (CDCl₃) δ 6.05 (dd, *J* = 6.0, 3.3 Hz, 1H), 5.84 (dd, *J* = 5.4, 1.8 Hz, 1H), 4.75 (m, 1H), 4.25 (m, 1H), 4.11 (m, 1H), 3.98 (m, 1H), 3.78 (br s, 3H), 2.58 (m, 1H), 2.30 (m, 2H).

***cis*-1-[4-(Methoxycarbonyloxymethyl)-2-cyclopenten-1-yl]-5-iodocytosine (**12**).** Tri-isopropyl phosphite (0.42 mL, 1.74 mmol) was added to a solution of Pd(OAc)₂ (0.098 g, 0.434 mmol) at ambient temperature in anhydrous THF (8.0 mL) under argon. After stirring this solution for 5 min, *n*-BuLi (1.6 *N* in hexane, 0.54 mL, 0.868 mmol) was added at ambient temperature. The resulting mixture was stirred for 5 min to obtain the tetrakis(triisopropylphosphite) palladium-(0) catalyst. The *in situ* prepared Pd(0) catalyst was added to a solution of 5-iodocytosine (**11**) (1.24 g, 5.2 mmol) and

sodium hydride (0.208 g, 5.2 mmol) in anhydrous DMSO (16.0 mL) *via* cannula at ambient temperature. Next, a solution of dicarbonate **10** (1.0 g, 4.34 mmol) in anhydrous THF (8.0 mL) was added to the reaction mixture. After stirring for 2 h, the reaction mixture was diluted with CHCl₃ (20 mL) and washed with a brine solution (20 mL $\times$ 3). The aqueous phase was extracted with CHCl₃ (20 mL $\times$ 2). The organic phase was collected, dried with anhydrous MgSO₄, and concentrated by rotary-evaporation. The residue was purified by silica-gel column chromatography (CHCl₃:MeOH = 20:1, v/v) to give a white solid **12** (*R*_f = 0.15; 1.19 g, 70%). mp 180 °C; IR (thin film): 3456, 3062, 2995, 2950, 1748, 1714, 1649, 1622, 1478, 1281, 957 cm⁻¹; ¹H NMR (DMSO-*d*₆) δ 7.69 (br s, 1H), 7.61 (s, 1H), 6.51 (br s, 1H), 6.05 (m, 1H), 5.79 (m, 1H), 5.50 (m, 1H), 4.21 - 4.07 (m, 2H), 3.70 (s, 3H), 3.03 (m, 1H), 2.60 (m, 1H), 1.30 (m, 1H); ¹³C NMR (DMSO-*d*₆) δ 164.10, 155.66, 154.97, 148.44, 137.65, 131.72, 69.90, 62.02, 56.70, 55.09, 44.30, 34.23; HRMS: calculated for C₁₂H₁₄IN₃O₄ [M⁺, FAB] 392.0107, found 392.0108.

***cis*-1-[4-(Methoxycarbonyloxymethyl)-2-cyclopenten-1-yl]-5-tributylstannylcytosine (**13**).** Bis(tributyltin) (0.4 mL, 0.75 mmol) and Pd₂(dba)₃ (0.023 g, 0.025 mmol) were added to a solution of compound **12** (0.20 g, 0.5 mmol) in anhydrous DMF (10 mL) at ambient temperature. The reaction mixture was stirred at 60 °C for 12 h. After removal of the solvent by rotary-evaporation, the residue was diluted with methanol (10 mL). Silica-gel (2.0 g) was added to the solution, and the resulting suspension was dried under reduced pressure. A pale yellow gummy solid **13** (*R*_f = 0.27; 0.155 g, 47%) was obtained from preloaded silica-gel column chromatography (CH₂Cl₂:MeOH = 20:1, v/v). IR (thin film): 3454, 3005, 2970, 2956, 2872, 1739, 1366, 1228, 1217, 528 cm⁻¹; ¹H NMR (CDCl₃) δ 7.11 (s, 1H), 6.08 (m, 1H), 5.83 (s, 1H), 5.74 (m, 1H), 4.18-4.06 (m, 2H), 3.77 (s, 3H), 3.10 (m, 1H), 2.82 (m, 1H), 1.54-1.46 (m, 6H), 1.36-1.29 (m, 7H), 1.13-1.02 (m, 6H), 0.93-0.88 (m, 9H); ¹³C NMR (CDCl₃) δ 169.76, 156.87, 155.74, 147.33, 136.78, 131.85, 100.90, 70.12, 61.46, 54.82, 44.02, 34.67, 28.93, 27.21, 13.66, 9.68; HRMS: calculated for C₂₄H₄₃N₃O₄Sn [M⁺, FAB] 556.2202, found 556.2200.

***cis*-1-[4-(Hydroxymethyl)-2-cyclopenten-1-yl]-5-iodocytosine (**14**).** Compound **12** (0.20 g, 0.511 mmol) was added to 0.5 *N* aqueous potassium carbonate (10.0 mL) and stirred at room temperature for 12 h. The reaction mixture was neutralized to pH 7-8 with 1.0 *N* HCl. After removal of the solvent by rotary-evaporation, the residue was diluted with methanol (10 mL). Silica-gel (2.0 g) was added to the solution, and the resulting suspension was dried under reduced pressure. A white solid **14** (*R*_f = 0.14; 0.155 g, 91%) was obtained from preloaded silica-gel column chromatography (CH₂Cl₂:MeOH = 20:1 to 15:1, v/v). mp 192 °C; IR (thin film): 3423, 2940, 2872, 1632, 1596, 1484 cm⁻¹; ¹H NMR (DMSO-*d*₆) δ 7.79 (s, 1H), 7.68 (br s, 1H), 6.48 (br s, 1H), 6.80 (m, 1H), 5.69 (m, 1H), 5.50 (m, 1H), 4.75 (m, 1H), 3.52-3.35 (m, 2H), 2.79 (m, 1H), 2.49 (m, 1H), 1.32 (m, 1H); ¹³C NMR (DMSO-*d*₆) δ 163.99, 155.00, 148.95, 139.97, 130.60, 63.69, 61.82, 56.31, 47.71, 34.03; HRMS: calculated

for $C_{10}H_{13}N_3O_2I$ [M^+ , FAB] 334.0053, found 334.0051.

***cis*-1-[4-(Hydroxymethyl)-2-cyclopenten-1-yl]-5-[^{124}I]iodocytosine (3).** Approximately 4.0 mg of the tributylstannylated compound **13** was dissolved in 5.0 mL of MeOH. The amount of radioiodide (0.3 mCi (11.1 MBq) $Na^{124}I$) in 0.01 *N* aqueous NaOH was added to the precursor **13** solution (100 μ L). The reaction mixture was acidified to a pH of 4.0 with 1.0 *N* aqueous HCl, and 30% H_2O_2 (50 μ L) was added. After stirring for 10 min at ambient temperature, the reaction mixture was quenched with saturated $NaHSO_3$. 1.0 *N* aqueous K_2CO_3 (100 μ L) was added. After stirring for 1 h at ambient temperature, the reaction mixture was neutralized with 1.0 *N* aqueous HCl to pH 7. Purification of the radioiodinated product **3** was performed by HPLC separation on a μ Bondapak C18 column (7.8 mm $\times$ 300 mm). The product was eluted at a flow rate of 2.0 mL/min with 0.1% trifluoroacetic acid in water/ethanol (9:1, v/v). The retention time of the product was 19.65 min. The radiochemical purity of the product was determined by the HPLC system used for purification and radio-thin layer chromatography (radio-TLC). Radio-TLC was performed on a silica-gel plate (Merck 60 F254 aluminum backed silica-gel plates), which was developed with acetonitrile/water (19:1, v/v) solvent (R_f = 0.23 for the product). The total elapsed time was about 110 minutes, including radioiodination and HPLC purification. The overall radiochemical yield and purity were more than 75% (8.33 MBq) and 95%, respectively.

***cis*-1-[4-(Methoxycarbonyloxymethyl)-2-cyclopenten-1-yl]-5-(2-trimethylsilylvinyl) cytosine (15).** (*E*)- $Bu_3SnCH=CHSiMe_3$ (2.0 g, 5.11 mmol) and $Pd_2(dba)_3$ (0.234 g, 0.256 mmol) were added to a solution of compound **12** (1.0 g, 2.56 mmol) in anhydrous DMF (30 mL) at ambient temperature. The reaction mixture was stirred at 60 $^{\circ}C$ for 12 h. After removal of the solvent by rotary-evaporation, the residue was diluted with methanol (10 mL). Silica-gel (~2.0 g) was added to the solution, and the resulting suspension was dried under reduced pressure. A yellow solid **15** (R_f = 0.23; 0.56 g, 60%) was obtained from preloaded silica-gel column chromatography (CH_2Cl_2 :MeOH = 20:1, v/v). mp 123 $^{\circ}C$; IR (thin film): 3428, 2956, 1750, 1646, 1474, 1444, 1273, 866, 841, 791 cm^{-1} ; 1H NMR ($CDCl_3$) δ 7.38 (s, 1H), 6.48 (d, J = 19.2 Hz, 1H), 6.13 (d, J = 19.2 Hz, 1H), 6.11 (m, 1H), 5.84 (m, 1H), 5.78 (m, 1H), 4.29–4.10 (m, 2H), 3.77 (s, 3H), 3.12 (m, 1H), 2.85 (m, 1H), 1.41 (m, 1H), 0.14 (m, 9H); ^{13}C NMR ($CDCl_3$) δ 163.45, 155.81, 155.80, 139.78, 137.45, 134.33, 131.91, 131.43, 107.24, 69.73, 61.85, 54.90, 44.23, 34.41, –1.26; HRMS: calculated for $C_{17}H_{25}N_3O_4Si$ [M^+ , FAB] 364.1693, found 364.1695.

***cis*-1-[4-(Hydroxymethyl)-2-cyclopenten-1-yl]-5-(2-trimethylsilylvinyl)cytosine (16).** Compound **15** (0.16 g, 0.44 mmol) was added to 0.5 *N* aqueous potassium carbonate (1.0 mL) and stirred at room temperature for 12 h. The reaction mixture was neutralized to pH 7–8 with 1.0 *N* HCl. After removal of the solvent by rotary-evaporation, the residue was diluted with methanol (10 mL). Silica-gel (~2.0 g) was added to the solution, and the resulting suspension was dried under reduced pressure. A yellow solid **16** (R_f = 0.25; 0.125

g, 93%) was obtained from preloaded silica-gel column chromatography ($CHCl_3$:MeOH = 15:1 to 10:1, v/v). mp 166 $^{\circ}C$; IR (thin film): 3341, 3202, 2953, 2895, 1644, 1592, 1507, 1475, 1248, 866, 840 cm^{-1} ; 1H NMR ($CDCl_3$) δ 7.55 (s, 1H), 6.44 (d, J = 19.2 Hz, 1H), 6.14 (m, 1H), 6.08 (d, J = 19.2 Hz, 1H), 5.74 (m, 1H), 5.73 (m, 1H), 4.77 (m, 1H), 3.77–3.66 (m, 2H), 2.79 (m, 1H), 2.75 (m, 1H), 1.50 (m, 1H), 0.13 (s, 9H); ^{13}C NMR ($CDCl_3$) δ 163.17, 155.88, 140.59, 138.67, 134.21, 131.95, 130.98, 106.89, 65.04, 62.48, 47.38, 34.07, –1.27; HRMS: calculated for $C_{15}H_{23}N_3O_2Si$ [M^+ , FAB] 306.1638, found 306.1642.

***cis*-1-[4-(Hydroxymethyl)-2-cyclopenten-1-yl]-5-(2-iodovinyl)cytosine (17).** ICl (0.053 g, 0.33 mmol) was added into solution of compound **16** (0.10 g, 0.33 mmol) in acetonitrile (8.0 mL). The reaction mixture was stirred at ambient temperature for 30 min. After removal of the solvent by rotary-evaporation, a yellow solid **17** (R_f = 0.24; 0.082 g, 70%) was obtained from silica-gel column chromatography (CH_2Cl_2 :MeOH = 10:1, v/v). mp 145 $^{\circ}C$; IR (thin film): 3362, 3255, 3050, 2929, 2873, 1635, 1501, 1488, 1401, 1290, 1223 cm^{-1} ; 1H NMR ($DMSO-d_6$) δ 7.68 (s, 1H), 7.19 (br s, 1H), 7.28 (d, J = 14.6 Hz, 1H), 6.63 (d, J = 14.6 Hz, 1H), 6.08 (m, 1H), 5.71 (m, 1H), 5.56 (m, 1H), 4.80 (m, 1H), 3.54–3.38 (m, 2H), 2.80 (m, 1H), 2.51 (m, 1H), 1.35 (m, 1H); ^{13}C NMR ($DMSO-d_6$) δ 162.44, 155.23, 140.36, 139.70, 136.22, 130.72, 105.48, 77.29, 63.81, 61.76, 47.75, 34.12; HRMS: calculated for $C_{12}H_{15}N_3O_2I$ [M^+ , FAB] 360.0209, found 360.0205.

***cis*-1-[4-(Hydroxymethyl)-2-cyclopenten-1-yl]-5-(2-[^{124}I]iodovinyl)cytosine (4).** Approximately 4.0 mg of trimethylsilyl compound **16** was dissolved in 5.0 mL of MeOH. The amount of radioiodide (0.3 mCi (11.1 MBq) $Na^{124}I$) in 0.01 *N* aqueous NaOH was added to the precursor **16** solution (100 μ L). The reaction mixture was acidified to a pH of 4.0 with 1.0 *N* aqueous HCl, and 30% H_2O_2 (50 μ L) was added. After stirring for 10 min at ambient temperature, the reaction mixture was quenched with saturated $NaHSO_3$. Purification of the radioiodinated product **4** was performed by HPLC separation on a μ Bondapak C18 column (7.8 mm $\times$ 300 mm). The product was eluted at a flow rate of 2.0 mL/min with 0.1% trifluoroacetic acid in water/acetonitrile (4:1, v/v). The retention time of the product was 17.0 min. The radiochemical purity of the product was determined by the HPLC system used for purification and radio-thin layer chromatography (radio-TLC). Radio-TLC was performed on a silica-gel plate (Merck 60 F254 aluminum backed silica-gel plates), which was developed with acetonitrile/water (19:1, v/v) solvent (R_f = 0.53 for the product). The total elapsed time was about 60 minutes, including radioiodination and HPLC purification. The overall radiochemical yield and purity were more than 80% (8.88 MBq) and 95%, respectively.

Acknowledgments. This research was supported by the Basic Science Research Program (2009-0070920) and the Nuclear R&D program (2010-0005313) through the National Research Foundation of Korea (NRF) funded by the Ministry of Education, Science and Technology. H.A. greatly appreciates

the Korean Ministry of Education through the second stage of the BK21 project for the Hanyang University graduate program and Seoul Metropolitan Government for Seoul Science Fellowship for financial support.

References

- (a) Kwon, H. C.; Kim, J. H.; Kim, K. C.; Lee, K. H.; Lee, J. H.; Lee, B. H.; Lee, K. H.; Jang, J. J.; Lee, C. T.; Lee, H.; Kim, C. M. *Mol. Cells* **2001**, *11*, 170. (b) Mullen, C. A. *Pharm. Ther.* **1994**, *63*, 199. (c) Choi, T. H.; Ahn, S. H.; Kwon, H. C.; Choi, C. W.; Awh, O. D.; Lim, S. M. *Appl. Radiat. Isot.* **2004**, *60*, 15. (c) Haberkorn, U.; Khazaie, K.; Morr, I.; Altmann, A.; Muller, M.; Kaick, G. V. *Nucl. Med. Biol.* **1998**, *25*, 367.
- (a) Tjuvajev, J. G.; Doubrovin, M.; Akhurst, T.; Cai, S.; Alauddin, J. M. M.; Finn, R.; Bornmann, W.; Thaler, H.; Conti, P. S.; Blasberg, R. G. *J. Nucl. Med.* **2002**, *43*, 1072. (b) Manger, T. J.; Klecker, R. W.; Anderson, L.; Shields, A. F. *Nucl. Med. Biol.* **2003**, *30*, 215. (c) Morin, K. W.; Atrazheva, E. D.; Knaus, E. E.; Wiebe, L. I. *J. Med. Chem.* **1997**, *40*, 2184. (d) Alauddin, M. M.; Shahinian, A.; Park, R.; Tohme, M.; Fissekis, J. D.; Conti, P. S. *J. Nucl. Med.* **2004**, *45*, 2063.
- (a) Morin, K. W.; Duan, W.; Zhou, A.; Xu, L.; Moharram, S.; Knaus, E. E.; McEwan, A. J. B.; Wiebe, L. I. *Nucl. Med. Biol.* **2004**, *31*, 623. (b) Toyohara, J.; Hayashi, A.; Sato, M.; Tanaka, H.; Haraguchi, K.; Yoshimura, Y.; Yonekura, Y.; Fujibayashi, Y. *J. Nucl. Med.* **2002**, *43*, 1218. (c) Toyohara, J.; Hayashi, A.; Sato, M.; Gogami, A.; Tanaka, H.; Haraguchi, K.; Yoshimura, Y.; Kumamoto, H.; Yonekura, Y.; Fujibayashi, Y. *Nucl. Med. Biol.* **2003**, *30*, 687.
- (a) Borthwick, A. D.; Biggadike, K. *Tetrahedron* **1992**, *48*, 571. (b) Herdewijn, P.; Clercq, E. D.; Balzarini, J.; Vanderhaeghe, H. *J. Med. Chem.* **1985**, *28*, 550. (c) Shi, J.; McAtee, J. J.; Wirtz, S. S.; Tharnish, P.; Juodawlkis, A.; Liotta, D. C.; Schinazi, R. F. *J. Med. Chem.* **1999**, *42*, 859. (d) Wang, J.; Jin, Y.; Rapp, K. L.; Benneth, M.; Schinazi, R. F.; Chu, C. K. *J. Med. Chem.* **2005**, *48*, 3736. (e) Agrofolio, L.; Suhas, E.; Farese, A.; Condom, R.; Challand, S. R.; Earl, R. A.; Guedj, R. *Tetrahedron* **1994**, *50*, 10611.
- (a) An, G.; Rhee, H. *Nucleosides Nucleotides Nucleic Acids* **2000**, *19*, 1111. (b) An, G.; Rhee, H. *Nucleosides Nucleotides Nucleic Acids* **2002**, *21*, 65. (c) An, G.; Rhee, H. *Nucleosides Nucleotides Nucleic Acids* **2003**, *22*, 437.
- Ahn, H.; Choi, T. H.; Castro, K. D. A.; Lee, K. C.; Kim, B.; Moon, B. S.; Hong, S. H.; Lee, J. C.; Chun, K. S.; Cheon, G. J.; Lim, S. M.; An, G.; Rhee, H. *J. Med. Chem.* **2007**, *50*, 6032.
- Kim, E.; Choi, T. H.; Ahn, S. H.; Kim, B.; Park, H.; Cheon, G. J.; Rhee, H.; An, G. *Nucl. Med. Mol. Imaging* **2009**, *43*, 478.
- Kim, J.; Castro, K. D. A.; Lim, M.; Rhee, H. *Tetrahedron* **2010**, *66*, 3995.
- (a) Schneekloth, J. S.; Mathieu, P.; Crews, C. M. *Eur. J. Org. Chem.* **2007**, *40*. (b) Negishi, E. *Handbook of Organopalladium Chemistry for Organic Synthesis*; John Wiley & Sons, Inc.: Hoboken, N. J., USA, **2002**, *1*, 693-719.
- (a) Sabbatini, P.; Wellendorph, P.; Hoeg, S.; Pedersen, M. H. F.; Brauner-Osborne, H.; Martiny, L.; Froelund, B.; Clausen, R. P. *J. Med. Chem.* **2010**, *53*, 6506. (b) Wang, B.; Qin, L.; Neumann, K. D.; Uppaluri, S.; Cerny, R. D.; Magno, S. G. D. *Org. Lett.* **2010**, *12*, 3352. (c) Watanabe, H.; Ono, M.; Haratake, M.; Kobashi, N.; Saji, H.; Nakayama, M. *Bioorg. Med. Chem.* **2010**, *18*, 4740.
- (a) Durka, K.; Gorka, J.; Kurach, P.; Lulinski, S.; Serwatowski, J. *J. Organomet. Chem.* **2010**, *695*, 2635. (b) Widhalm, M.; Abraham, M.; Arion, V. B.; Saarsalu, S.; Maeorg, U. *Tetrahedron: Asymm.* **2010**, *21*, 1971. (c) Denmark S. E.; Muhuhi, J. M. *J. Am. Chem. Soc.* **2010**, *132*, 11768.
- (a) Bukovec, C.; Wesquet, A. O.; Kazmaier, U. *Eur. J. Org. Chem.* **2011**, 1047. (b) Perez-Temprano, M. H.; Gallego, A. M.; Casares, J. A.; Espinet, P. *Organometallics* **2011**, *30*, 611. (c) Yabe, Y.; Maegawa, T.; Monguchi, Y.; Sajiki, H. *Tetrahedron* **2010**, *66*, 8654.

Supporting Information

Synthesis of Radioiodinated Carbocyclic Cytosine Analogues

Hyunseok Ahn,[†] Gwangil An,[‡] and Hakjune Rhee^{†,§,*}

[†]*Department of Bionanotechnology, Hanyang University, Kyunggi-do 426-791 Korea. *E-mail: hrhee@hanyang.ac.kr*

[‡]*Laboratory of Radiopharmaceuticals and Laboratory of Nuclear Medicine, Seoul 139-706, Korea*

[§]*Department of Chemistry and Applied Chemistry, Hanyang University, Kyunggi-do 426-791 Korea*

Received March 29, 2011, Accepted April 21, 2011

cis-1-[4-(Methoxycarbonyloxymethyl)-2-cyclopenten-1-yl]-5-iodocytosine (12)

ahs-269-d41c-coupling
Pulse Sequence: s2pul
Solvent: DMSO
Temp. 25.0 C / 298.1 K
File: ahs-269-DMSO-1H
Mercury-300 "nmr300"

Relax. delay 2.000 sec
Pulse 85.5 degrees
Acq. time 1.999 sec
Width 4001.6 Hz
16 repetitions
OBSERVE H1 300.0746498 MHz
DATA PROCESSING
FT size 16384
Total time 1 min, 5 sec

[Mass Spectrum]
Data : #45269-C12H14IN3O4 Date : 30-Nov-2010 14:59
Sample : -
Note : -
Inlet : Direct Ion Mode : FAB+
Spectrum Type : Normal Ion (EF-Linear)
RT : 0.38 min Scan# : (8,9)
BP : m/z 430.1736 Int. : 99.88
Output m/z range : 388.4710 to 404.9018 Cut Level : 0.00 %

***cis*-1-[4-(Methoxycarbonyloxymethyl)-2-cyclopenten-1-yl]-5-tributylstannylcytosine (13)**

[Mass Spectrum]
Data : FHS276-C18H12IN302 Date : 30-Nov-2010 14:49
Sample : -
Note : -
Inlet : Direct Ion Mode : FRB+
Spectrum Type : Normal Ion (EF-Linear)
RT : 0.33 min Scan# : (7,8)
BP : m/z 307.1035 Int. : 91.61
Output m/z range : 325.7889 to 346.6213 Cut Level : 0.00 %

cis-1-[4-(Hydroxymethyl)-2-cyclopenten-1-yl]-5-iodocytosine (14)

Std proton

Sample: ahs-271-1H
 File: /home/vnmr1/vnmrsys/data/hank/20100205-ahs-271-1H.fid
 Pulse Sequence: s2pu1
 Solvent: cdcl3
 Ambient temperature
 Operator: vnmr1
 File: 20100205-ahs-271-1H
 VNMR-S500 "hyunmr600"

Relax. delay 1.000 sec
 Pulse 45.0 degrees
 Acq. time 2.049 sec
 Width 8012.8 Hz
 16 repetitions
 OBSERVE H1, 499.9635234 MHz
 DATA PROCESSING
 Resol. enhancement -0.0 Hz
 FT size 65536
 Total time 0 min, 55 sec

[Mass Spectrum]
Data : FHS271-C24H43N3O4Sn Date : 30-Nov-2010 14:38
Sample : -
Note : -
Inlet : Direct Ion Mode : FAB+
Spectrum Type : Normal Ion [EF-Linear]
RT : 0.68 min Scan# : (14,15)
BP : m/z 556.2200 Int. : 72.35
Output m/z range : 543.6202 to 568.3680 Cut Level : 0.00 %
1605761

***cis*-1-[4-(Methoxycarbonyloxymethyl)-2-cyclopenten-1-yl]-5-(2-trimethylsilylvinyl)cytosine (15)**

[Mass Spectrum]
Data : F45275-C17H25N3O4S1 Date : 30-Nov-2010 14:33
Sample: -
Note: -
Inlet : Direct Ion Mode : FAB+
Spectrum Type : Normal Ion (EF-Linear)
RT : 0.28 min Scan# : (6,7)
BP : m/z 364.1695 Int. : 59.98
Output m/z range : 349.1395 to 380.8308 Cut Level : 0.00 %

cis-1-[4-(Hydroxymethyl)-2-cyclopenten-1-yl]-5-(2-trimethylsilylvinyl)cytosine (16)

[Mass Spectrum]
Data : R45279-C15H23N3O2S1 Date : 30-Nov-2010 14:28
Sample : -
Note : -
Inlet : Direct Ion Mode : FAB+
Spectrum Type : Normal Ion [EF-Linear]
RT : 0.28 min Scan# : (6,7)
BP : m/z 306.1642 Int. : 94.58
Output m/z range : 295.4970 to 318.1306 Cut Level : 0.00 %

***cis*-1-[4-(Hydroxymethyl)-2-cyclopenten-1-yl]-5-(2-iodovinyl)cytosine (17)**

[Mass Spectrum]
Data : FHS280-d4IVC-C12H14IN3O2 Date : 16-Dec-2010 16:56
Sample : -
Note : -
Inlet : Direct Ion Mode : FAB+
Spectrum Type : Normal Ion [EF-Linear]
RT : 0.33 min Scan# : (7,8)
BP : m/z 391.2881 Int. : 75.68
Output m/z range : 346.2908 to 377.6261 Cut Level : 0.00 %
1217331

Radio-TLC and HPLC Analysis for Radioiodination of Carbocyclic d4IC (3)

Radio-TLC analysis

HPLC analysis

Radio-TLC and HPLC Analysis for Radioiodination of Carbocyclic d4IVC (4)

Radio-TLC analysis

HPLC analysis