JACOBI OPERATORS ALONG THE STRUCTURE FLOW ON REAL HYPERSURFACES IN A NONFLAT COMPLEX SPACE FORM II

U-HANG KI AND HIROYUKI KURIHARA

ABSTRACT. Let M be a real hypersurface of a complex space form with almost contact metric structure (ϕ, ξ, η, g) . In this paper, we study real hypersurfaces in a complex space form whose structure Jacobi operator $R_\xi = R(\cdot, \xi)\xi$ is ξ -parallel. In particular, we prove that the condition $\nabla_\xi R_\xi = 0$ characterizes the homogeneous real hypersurfaces of type A in a complex projective space or a complex hyperbolic space when $R_\xi \phi S = R_\xi S \phi$ holds on M , where S denotes the Ricci tensor of type (1,1) on M .

1. Introduction

Let $(M_n(c), J, \tilde{g})$ be a complex n -dimensional complex space form with Kähler structure $(J, \tilde{g})$ of constant holomorphic sectional curvature $4c$ and let M be an orientable real hypersurface in $M_n(c)$. Then M has an almost contact metric structure (ϕ, ξ, η, g) induced from $(J, \tilde{g})$.

Typical examples of real hypersurfaces in $M_n(c)$, $c \neq 0$, are homogeneous ones. Takagi [12], [13] classified homogeneous real hypersurfaces of a complex projective space $P_n\mathbb{C}$ into six model spaces. On the other hand, Cecil and Ryan [2] extensively studied a Hopf hypersurface, which is realized as tubes over certain submanifolds in $P_n\mathbb{C}$, by using its focal map. By making use of those results and the mentioned work of Takagi, Kimura [7] proved the local classification theorem for Hopf hypersurfaces of $P_n\mathbb{C}$ all of whose principal curvatures are constants. For the case where a complex hyperbolic space $H_n\mathbb{C}$, Berndt [1] proved the classification theorem for Hopf hypersurfaces whose all principal curvatures are constants. Among the several types of real hypersurfaces appeared in model spaces ([1], [12]), a particular type of tubes over totally geodesic $P_k\mathbb{C}$ or $H_k\mathbb{C}$ ($0 \leq k \leq n-1$) adding a horosphere in $H_n\mathbb{C}$, which is called type A , has a lot of nice geometric properties. For example, Okumura [9] (resp. Montiel and Romero [8]) showed that a real hypersurface in $P_n\mathbb{C}$ (resp.

Received August 17, 2010; Revised June 1, 2011.

2010 *Mathematics Subject Classification.* Primary 53B20, 53C15, 53C25.

Key words and phrases. complex space form, real hypersurface, structure Jacobi operator, Ricci tensor.

$H_n\mathbb{C}$) is locally congruent to one of real hypersurfaces of type A if and only if the Reeb flow ξ is isometric or equivalently the structure operator ϕ commutes with the shape operator H .

The structure Jacobi operator $R_\xi = R(\cdot, \xi)\xi$ has a fundamental role in contact geometry. Ortega, Pérez and Santos [10] have proved that there are no real hypersurfaces in $P_n\mathbb{C}$, $n \geq 3$ with parallel structure Jacobi operator $\nabla R_\xi = 0$. More generally, such a result has been extended by [11] due to them. In this paper, motivated by results mentioned above we consider the parallelism of the structure Jacobi operator R_ξ in the direction of the structure vector field, that is $\nabla_\xi R_\xi = 0$. Recently we have some classification theorems for real hypersurfaces in a nonflat complex space form with respect to the parallelism of R_ξ (cf. [3], [4], [5]).

The main purpose of this paper is to classify real hypersurfaces in a nonflat complex space form $M_n(c)$ which satisfies $\nabla_\xi R_\xi = 0$ and at the same time $R_\xi\phi S = R_\xi S\phi$, where S denotes the Ricci tensor of the hypersurface. Our main result is as follows:

Main Theorem. *Let M be a real hypersurface in a complex space form $M_n(c)$, $c \neq 0$, $n \geq 3$ which satisfies $\nabla_\xi R_\xi = 0$. Then M holds $R_\xi\phi S = R_\xi S\phi$ if and only if $H\xi = 0$ or M is locally congruent to one of the following:*

- (I) *in case that $M_n(c) = P_n\mathbb{C}$,*
 - (A₁) *a geodesic hypersphere of radius r , where $0 < r < \pi/2$ and $r \neq \pi/4$,*
 - (A₂) *a tube of radius r over a totally geodesic $P_k\mathbb{C}$ for some $k \in \{1, \dots, n-2\}$, where $0 < r < \pi/2$ and $r \neq \pi/4$;*
- (II) *in case that $M_n(c) = H_n\mathbb{C}$,*
 - (A₀) *a horosphere,*
 - (A₁) *a geodesic hypersphere or a tube over a complex hyperbolic hyperplane $H_{n-1}\mathbb{C}$,*
 - (A₂) *a tube over a totally geodesic $H_k\mathbb{C}$ for some $k \in \{1, \dots, n-2\}$.*

We note that every Hopf hypersurface always satisfies condition $R_\xi\phi S = R_\xi S\phi$.

All manifolds in this paper are assumed to be connected and of class C^∞ and the real hypersurfaces are supposed to be oriented.

2. Preliminaries

We denote by $M_n(c)$, $c \neq 0$, be a nonflat complex space form with the Fubini-Study metric $\tilde{g}$ of constant holomorphic sectional curvature $4c$ and Levi-Civita connection $\tilde{\nabla}$. For an immersed $(2n-1)$ -dimensional Riemannian manifold $\tau : M \rightarrow M_n(c)$, the Levi-Civita connection ∇ of induced metric and the shape operator H of the immersion are characterized

$$\tilde{\nabla}_X Y = \nabla_X Y + g(HX, Y)\nu, \quad \tilde{\nabla}_X \nu = -HX$$

for any vector fields X and Y on M , where g denotes the Riemannian metric of M induced from $\tilde{g}$ and ν a unit normal vector on M . In the sequel the indices $i, j, k, l, \dots$ run over the range $\{1, 2, \dots, 2n-1\}$ unless otherwise stated. For a local orthonormal frame field $\{e_i\}$ of M , we denote the dual 1-forms by $\{\theta_i\}$. Then the connection forms θ_{ij} are defined by

$$d\theta_i + \sum_j \theta_{ij} \wedge \theta_j = 0, \quad \theta_{ij} + \theta_{ji} = 0.$$

Then we have

$$\nabla_{e_i} e_j = \sum_k \theta_{kj}(e_i) e_k = \sum_k \Gamma_{kij} e_k,$$

where we put $\theta_{ij} = \sum_k \Gamma_{ijk} \theta_k$. The almost contact metric structure $(\phi = (\phi_{ij}), \xi = \sum_i \xi_i e_i)$ is induced on M by following equation:

$$J(e_i) = \sum_j \phi_{ji} e_j + \xi_i \nu.$$

The structure tensor ϕ and the structure vector ξ satisfy

$$\begin{aligned} \sum_k \phi_{ik} \phi_{kj} &= \xi_i \xi_j - \delta_{ij}, \quad \sum_j \xi_j \phi_{ij} = 0, \quad \sum_i \xi_i^2 = 1, \quad \phi_{ij} + \phi_{ji} = 0, \\ (2.1) \quad d\phi_{ij} &= \sum_k (\phi_{ik} \theta_{kj} - \phi_{jk} \theta_{ki} - \xi_i h_{jk} \theta_k + \xi_j h_{ik} \theta_k), \\ d\xi_i &= \sum_j \xi_j \theta_{ji} - \sum_{j,k} \phi_{ji} h_{jk} \theta_k. \end{aligned}$$

We denote the components of the shape operator or the second fundamental tensor H of M by h_{ij} . The components $h_{ij;k}$ of the covariant derivative of H are given by $\sum_k h_{ij;k} \theta_k = dh_{ij} - \sum_k h_{ik} \theta_{kj} - \sum_k h_{jk} \theta_{ki}$. Then we have the equation of Gauss and Codazzi

$$(2.2) \quad R_{ijkl} = c(\delta_{ik} \delta_{jl} - \delta_{il} \delta_{jk} + \phi_{ik} \phi_{jl} - \phi_{il} \phi_{jk} + 2\phi_{ij} \phi_{kl}) + h_{ik} h_{jl} - h_{il} h_{jk},$$

$$(2.3) \quad h_{ij;k} - h_{ik;j} = c(\xi_k \phi_{ij} + \xi_i \phi_{kj} - \xi_j \phi_{ik} - \xi_i \phi_{jk}),$$

respectively.

From (2.2) the structure Jacobi operator $R_\xi = (\Xi_{ij})$ is given by

$$(2.4) \quad \Xi_{ij} = \sum_{k,l} h_{ik} h_{jl} \xi_k \xi_l - \sum_{k,l} h_{ij} h_{kl} \xi_k \xi_l + c \xi_i \xi_j - c \delta_{ij}.$$

From (2.2) the Ricci tensor $S = (S_{ij})$ is given by

$$(2.5) \quad S_{ij} = (2n+1)c\delta_{ij} - 3c\xi_i \xi_j + h h_{ij} - \sum_k h_{ik} h_{kj},$$

where $h = \sum_i h_{ii}$.

First we remark:

Lemma 1 ([5]). *Let U be an open set in M and F a smooth function on U . We put $dF = \sum_i F_i \theta_i$. Then we have*

$$F_{ij} - F_{ji} = \sum_k F_k \Gamma_{kij} - \sum_k F_k \Gamma_{kji}.$$

Now we retake a local orthonormal frame field e_i in such a way that

- $e_1 = \xi$,
- e_2 is in the direction of $\sum_{i=2}^{2n-1} h_{1i} e_i$,
- $e_3 = \phi e_2$.

Then we have

$$(2.6) \quad \xi_1 = 1, \quad \xi_i = 0 \quad (i \geq 2), \quad h_{1j} = 0 \quad (j \geq 3) \quad \text{and} \quad \phi_{32} = 1.$$

We put $\alpha := h_{11}, \beta := h_{12}, \gamma := h_{22}, \varepsilon := h_{23}$ and $\delta := h_{33}$.

Hereafter the indices $p, q, r, s, \dots$ run over the range $\{4, 5, \dots, 2n-1\}$ unless otherwise stated.

Since $d\xi_i = 0$, we have

$$(2.7) \quad \begin{aligned} \theta_{12} &= \varepsilon \theta_2 + \delta \theta_3 + \sum_p h_{3p} \theta_p, \\ \theta_{13} &= -\beta \theta_1 - \gamma \theta_2 - \varepsilon \theta_3 - \sum_p h_{2p} \theta_p, \\ \theta_{1p} &= \sum_q \phi_{qp} h_{q2} \theta_2 + \sum_q \phi_{qp} h_{q3} \theta_3 + \sum_{q,r} \phi_{qp} h_{qr} \theta_r. \end{aligned}$$

We put

$$(2.8) \quad \theta_{23} = \sum_i X_i \theta_i, \quad \theta_{2p} = \sum_i Y_{pi} \theta_i, \quad \theta_{3p} = \sum_i Z_{pi} \theta_i.$$

Then it follows from $d\phi_{2i} = 0$ that $Y_{pi} = -\sum_q \phi_{pq} Z_{qi}$ or $Z_{pi} = \sum_q \phi_{pq} Y_{qi}$. The equations (2.4) and (2.5) are rewritten as

$$(2.9) \quad \Xi_{ij} = -\alpha h_{ij} + h_{1i} h_{1j} + c \delta_{i1} \delta_{j1} - c \delta_{ij},$$

$$(2.10) \quad S_{ij} = h h_{ij} - \sum_k h_{ik} h_{jk} - 3c \delta_{i1} \delta_{j1} + (2n+1) c \delta_{ij},$$

respectively.

3. Real hypersurfaces satisfying $\nabla_\xi R_\xi = 0$ and $R_\xi \phi S = R_\xi S \phi$

First we assume that $\nabla_\xi R_\xi = 0$. The components $\Xi_{ij;k}$ of the covariant derivativation of $R_\xi = (\Xi_{ij})$ is given by

$$\sum_k \Xi_{ij;k} \theta_k = d\Xi_{ij} - \sum_k \Xi_{kj} \theta_{ki} - \sum_k \Xi_{ik} \theta_{kj}.$$

Substituting (2.9) into the above equation we have

$$\begin{aligned}
 \sum_k \Xi_{ij;k} \theta_k &= -(d\alpha)h_{ij} - \alpha dh_{ij} + (dh_{1i})h_{1j} + h_{1i}(dh_{1j}) \\
 (3.1) \quad &+ \alpha \sum_k h_{kj} \theta_{ki} - \alpha h_{1j} \theta_{1i} - \beta h_{1j} \theta_{2i} - c \delta_{j1} \theta_{1i} \\
 &+ \alpha \sum_k h_{ik} \theta_{kj} - \alpha h_{1i} \theta_{1j} - \beta h_{1i} \theta_{2j} - c \delta_{i1} \theta_{1j}.
 \end{aligned}$$

In the following, we assume that $\beta \neq 0$.

Our assumption $\nabla_\xi R_\xi = 0$ is equivalent to $\Xi_{ij;1} = 0$, which can be stated as follows:

$$(3.2) \quad \varepsilon = 0, \quad \alpha\delta + c = 0, \quad h_{3p} = 0,$$

$$(3.3) \quad (\beta^2 - \alpha\gamma)_1 - 2\alpha \sum_p h_{2p} Y_{p1} = 0,$$

$$(3.4) \quad (\beta^2 - \alpha\gamma - c)X_1 + \alpha \sum_p h_{2p} Z_{p1} = 0,$$

$$(3.5) \quad (\alpha h_{2p})_1 + \alpha \sum_q h_{pq} Y_{q1} + (\beta^2 - \alpha\gamma)Y_{p1} - \alpha \sum_q h_{2q} \Gamma_{qp1} = 0,$$

$$(3.6) \quad \alpha h_{2p} X_1 - \sum_q (\alpha h_{qp} + c \delta_{pq}) Z_{q1} = 0,$$

$$(3.7) \quad (\alpha h_{pq})_1 - \alpha h_{2q} Y_{p1} - \alpha \sum_r h_{rq} \Gamma_{rp1} - \alpha h_{2p} Y_{q1} - \alpha \sum_r h_{pr} \Gamma_{rq1} = 0.$$

Hereafter we shall use (3.2) without quoting.

Furthermore we assume that $R_\xi \phi S = R_\xi S \phi$. Under the assumption $\nabla_\xi R_\xi = 0$, we have the following additional equations

$$(3.8) \quad \beta^2 - \alpha\gamma - c = 0,$$

$$(3.9) \quad \tilde{R}_\xi \tilde{\phi} A = 0,$$

$$(3.10) \quad \tilde{R}_\xi \tilde{\phi} \tilde{S} = \tilde{R}_\xi \tilde{S} \tilde{\phi},$$

where $A = {}^t(h_{24}, h_{25}, \dots, h_{2,2n-1})$, $\tilde{R}_\xi = (\Xi_{pq})$, $\tilde{\phi} = (\phi_{pq})$, $\tilde{S} = (S_{pq})$. From (3.2) and (3.8) we note

$$(3.11) \quad \delta \neq \gamma.$$

Now, properly speaking, we denote the equation (2.3) by (ijk) simply. Then we have the following equations (112)–(q3p).

$$(112) \quad \alpha_2 - \beta_1 = 0,$$

$$(212) \quad \beta_2 - \gamma_1 - 2 \sum_p h_{2p} Y_{p1} = 0,$$

$$(312) \quad (\alpha - \delta)\gamma - \beta X_2 + (\gamma - \delta)X_1 - \beta^2 - \sum_p h_{2p}Z_{p1} = -c,$$

$$(113) \quad \alpha_3 + 3\beta\delta - \alpha\beta + \beta X_1 = 0,$$

$$(213) \quad \beta_3 + \gamma\delta + (\gamma - \delta)X_1 - \beta^2 - \sum_p h_{2p}Z_{p1} = 0,$$

$$(313) \quad \beta X_3 + \delta_1 = 0,$$

$$(223) \quad \gamma_3 - 2\beta\delta + 2 \sum_p h_{2p}Y_{p3} + (\gamma - \delta)X_2 - \beta\gamma - \sum_p h_{2p}Z_{p2} = 0,$$

$$(323) \quad \sum_p h_{2p}Z_{p3} - \delta_2 - (\gamma - \delta)X_3 = 0,$$

$$(1p1) \quad \alpha_p + \beta Y_{p1} = 0,$$

$$(12p) \quad \beta_p + 2 \sum_{q,r} h_{2q}\phi_{rq}h_{rp} + \beta Y_{p2} + \alpha \sum_q \phi_{qp}h_{2q} = 0,$$

$$(13p) \quad 2\delta h_{2p} - \beta Y_{p3} - \alpha h_{2p} + \beta X_p = 0,$$

$$(22p) \quad \gamma_p + 2 \sum_q h_{2q}Y_{qp} - h_{2p2} - \sum_q h_{qp}Y_{q2} + \beta \sum_q \phi_{qp}h_{2q} + \gamma Y_{p2} \\ + \sum_q h_{2q}\Gamma_{qp2} = 0,$$

$$(23p) \quad \delta X_p + \beta h_{2p} - \gamma X_p + \sum_q h_{2q}Z_{qp} - h_{2p3} - \sum_q h_{qp}Y_{q3} \\ + \gamma Y_{p3} + \sum_q h_{2q}\Gamma_{qp3} = 0,$$

$$(33p) \quad \delta_p + h_{2p}X_3 - \sum_q h_{qp}Z_{q3} + \delta Z_{p3} = 0,$$

$$(21p) \quad \beta_p + \sum_{q,r} h_{2q}\phi_{rq}h_{rp} - h_{2p1} - \sum_q h_{qp}Y_{q1} + \gamma Y_{p1} + \sum_q h_{2q}\Gamma_{qp1} = 0,$$

$$(31p) \quad \delta h_{2p} - \alpha h_{2p} + \beta X_p - h_{2p}X_1 + \sum_q h_{qp}Z_{q1} - \delta Z_{p1} = 0,$$

$$(32p) \quad \delta X_p + \beta h_{2p} - \gamma X_p + \sum_q h_{2q}Z_{qp} + h_{2p}X_2 - \sum_q h_{pq}Z_{q2} + \delta Z_{p2} = 0,$$

$$(1pq) \quad 2 \sum_{r,s} h_{rp}\phi_{sr}h_{sq} - \alpha \sum_r \phi_{rp}h_{rq} + \alpha \sum_r \phi_{rq}h_{rp} \\ - \beta Y_{pq} + \beta Y_{qp} = -2c\phi_{pq},$$

$$(2pq) \quad h_{2pq} + \sum_r h_{rp}Y_{rq} - \beta \sum_r \phi_{rp}h_{rq} - \gamma Y_{pq} - \sum_r h_{2r}\Gamma_{rpq} - h_{2qp}$$

$$\begin{aligned}
& - \sum_r h_{rq} Y_{rp} + \beta \sum_r \phi_{rq} h_{rp} + \gamma Y_{qp} + \sum_r h_{2r} \Gamma_{rqp} = 0, \\
(q1p) \quad & \sum_{r,s} h_{rq} \phi_{sr} h_{sp} - \alpha \sum_r \phi_{rq} h_{rp} - \beta Y_{qp} - h_{pq1} \\
& + h_{2q} Y_{p1} + \sum_r h_{rq} \Gamma_{rp1} + h_{2p} Y_{q1} + \sum_r h_{rp} \Gamma_{rq1} = c \phi_{pq}, \\
(q3p) \quad & h_{3qp} - \varepsilon Y_{qp} - \delta Z_{qp} - \sum_r h_{3r} \Gamma_{rqp} - h_{2q} X_p + \sum_r h_{qr} Z_{rp} - h_{qp3} \\
& + h_{q2} Y_{p3} + h_{q3} Z_{p3} + \sum_r h_{qr} \Gamma_{rp3} + h_{p2} Y_{q3} + h_{p3} Z_{q3} \\
& + \sum_r h_{pr} \Gamma_{rq3} = 0.
\end{aligned}$$

Remark 1. We did not write $(p2q)$, $(3pq)$ and (pqr) since we need not use them.

4. Formulas and lemmas

In this section we study the crucial case where $\beta \neq 0$. By (3.6) and (31p) we have

$$(4.1) \quad \beta X_p = (\alpha - \delta) h_{2p}.$$

This and (13p) imply that

$$(4.2) \quad \beta Y_{p3} = \delta h_{2p}.$$

The equation (3.9) can be rewritten as

$$(4.3) \quad \sum_{q,r} (\alpha h_{pq} + c \delta_{pq}) \phi_{qr} h_{r2} = 0,$$

which, together with (4.2), implies

$$\beta \sum_q (h_{pq} - \delta \delta_{pq}) Z_{q3} = \delta \sum_{q,r} (h_{pq} - \delta \delta_{pq}) \phi_{qr} Y_{r3} = 0.$$

Hence it follows from (33p) and (1p1) that

$$(4.4) \quad \delta_p = -h_{2p} X_3 \quad \text{and} \quad \alpha_p = -\beta Y_{p1}.$$

Thus since (4.4) and $\alpha_p \delta + \alpha \delta_p = 0$ obtained from (3.2) we have

$$(4.5) \quad \beta \delta Y_{p1} = -\alpha h_{2p} X_3,$$

and so $\sum_p h_{2p} Z_{p1} = 0$. By (4.2), we have

$$(4.6) \quad \sum_p h_{2p} Z_{p3} = \sum_{p,q} h_{2p} \phi_{pq} Y_{q3} = \frac{\delta}{\beta} \sum_{p,q} h_{2p} \phi_{pq} h_{2q} = 0.$$

From (3.6), (4.3) and (4.5) we have

$$(4.7) \quad h_{2p}X_1 = 0.$$

Now we shall prove the following key lemma.

Lemma 2. $H(e_2) \in \text{span}\{e_1, e_2\}$.

Proof. Suppose that $h_{2p} \neq 0$. Then from (4.7) we have $X_1 = 0$. We can select the vector e_4 so that $h_{24} \neq 0$ and $h_{25} = \cdots = h_{2,2n-1} = 0$. We put $e_5 := \phi e_4$ and $\rho := h_{24}(\neq 0)$. Note that $\phi_{54} = 1$. Then by (4.3) we have

$$h_{55} = \delta, \quad h_{p5} = 0 \quad (p \neq 5).$$

Put $p = 5$ in (32p). Then since above equation and (4.1) we have $X_5 = 0$ and so $Z_{45} = 0$. Thus we have $Y_{55} = 0$. Furthermore, put $p = q = 5$ in (q1p). Then, since $\Gamma_{551} = Y_{55} = 0$, we have

$$(4.8) \quad \alpha_1 = \delta_1 = 0.$$

Thus, from (313), (323), (4.6) and (112) we have

$$(4.9) \quad X_3 = 0,$$

$$(4.10) \quad \alpha_2 = \delta_2 = 0,$$

$$(4.11) \quad \beta_1 = 0.$$

By (4.4) and (4.9) we have $\alpha_p = \delta_p = 0$. Thus it follows from (1p1) that

$$(4.12) \quad \alpha_p = \delta_p = Y_{p1} = Z_{p1} = 0.$$

Now we put $F = \alpha$, $i = 1$ and $j = p$ in Lemma 1. Then, from (2.7), (4.8), (4.10) and (4.12) we have

$$0 = \alpha_{1p} - \alpha_{p1} = \sum_k \alpha_k \Gamma_{k1p} - \sum_k \alpha_k \Gamma_{kp1} = \alpha_3(\Gamma_{31p} - \Gamma_{3p1}) = \alpha_3 h_{2p}.$$

Thus we have $\alpha_3 = 0$. Hence it follows from (4.8), (4.10) and (4.12) that α and δ are constants, which, together with (113), imply

$$(4.13) \quad \alpha = 3\delta.$$

On the other hand, taking account of the coefficient of $\theta_1 \wedge \theta_3$ in the exterior derivative of θ_{23} , we have $X_2 = -2\beta$. Thus, from (312) and (4.13) we have

$$2\delta\gamma + \beta^2 = -c.$$

Hence it follows from (3.8) that β and γ are constants. From (3.5) and (4.12) we have

$$h_{2p1} - \sum_q h_{2q} \Gamma_{qp1} = 0.$$

This, together with (21p), implies $\rho\delta = 0$, which is a contradiction. $\square$

Remark 2. Lemma 2 has been already proved in [5]. However in the case of our condition by using the equation (3.8) we have a short proof.

Owing to Lemma 2 the matrix (h_{pq}) is diagonalizable, that is, for a suitable choice of a orthonormal frame field $\{e_p\}$ we can set

$$h_{pq} = \lambda_p \delta_{pq}.$$

Then it is easy to see

$$(4.14) \quad \begin{aligned} \tilde{R}_\xi &= -((\alpha\lambda_p + c)\delta_{pq}), \\ \tilde{S} &= ((h\lambda_p - (\lambda_p)^2 + (2n+1)c)\delta_{pq}). \end{aligned}$$

Here we shall sum up all equations obtained from Lemma 2.

From (4.1), (4.2) and (4.4) we have

$$(4.15) \quad X_p = Y_{p1} = Z_{p1} = Y_{p3} = Z_{p3} = 0, \quad \alpha_p = \delta_p = 0.$$

Put $p = q$ in (3.7). Then we have

$$(4.16) \quad (\alpha\lambda_p)_1 = 0.$$

Moreover, from (112)–(32p) we have

$$(4.17) \quad \alpha_2 - \beta_1 = 0,$$

$$(4.18) \quad \beta_2 - \gamma_1 = 0,$$

$$(4.19) \quad \gamma\delta + \beta X_2 - (\gamma - \delta)X_1 = 0,$$

$$(4.20) \quad \alpha_3 + 3\beta\delta - \alpha\beta + \beta X_1 = 0,$$

$$(4.21) \quad \beta_3 - \alpha\delta + \gamma\delta + (\gamma - \delta)X_1 - \beta^2 = c,$$

$$(4.22) \quad \delta_1 + \beta X_3 = 0,$$

$$(4.23) \quad \gamma_3 - 2\beta\delta + (\gamma - \delta)X_2 - \beta\gamma = 0,$$

$$(4.24) \quad \delta_2 + (\gamma - \delta)X_3 = 0,$$

$$(4.25) \quad \beta_p = 0,$$

$$(4.26) \quad Y_{p2} = 0, \quad Z_{p2} = 0,$$

$$(4.27) \quad \gamma_p = 0.$$

It follows from (q1p) and (3.7) that

$$(4.28) \quad \alpha\beta Y_{qp} = \alpha\lambda_p\lambda_q\phi_{pq} - \alpha^2\lambda_p\phi_{pq} + \alpha_1\lambda_p\delta_{pq} - c\alpha\phi_{pq}.$$

From this, (2pq) and (q3p) we have

$$(4.29) \quad \begin{aligned} \beta^2(\lambda_p + \lambda_q)\phi_{pq} - (\lambda_p - \gamma)(\lambda_p\lambda_q - \alpha\lambda_q - c)\phi_{pq} \\ - (\lambda_q - \gamma)(\lambda_p\lambda_q - \alpha\lambda_p - c)\phi_{pq} = 0, \end{aligned}$$

$$(4.30) \quad (\lambda_q - \delta)(\alpha((\lambda_q)^2 - \alpha\lambda_q - c)\delta_{pq} + \alpha_1\lambda_q\phi_{pq}) - \alpha\beta(h_{qp3} + (\lambda_p - \lambda_q)\Gamma_{qp3}) = 0.$$

If $p = q$ in above equation, then we have

$$(4.31) \quad (\lambda_p - \delta)((\lambda_p)^2 - \alpha\lambda_p - c) - \beta(\lambda_p)_3 = 0.$$

5. Proof of Main Theorem

In this section we prove Main Theorem.

We first prove $\beta = 0$. Suppose that $\beta \neq 0$. It follows from (4.22) that (3.10) is equivalent to

$$\rho_p(\sigma_q - \sigma_p)\phi_{pq} = 0,$$

where $\rho_p = \alpha\lambda_p + c$, $\sigma_p = h\lambda_p - (\lambda_p)^2 + (2n+1)c$. Therefore if $\phi_{qp} \neq 0$, then we have

$$(5.1) \quad (\lambda_p - \lambda_q)(\alpha\lambda_p + c)(h - \lambda_q - \lambda_p) = 0.$$

Here we assert that if $\phi_{pq} \neq 0$, then $\lambda_p = \lambda_q$. To prove this, we assume that there exist indices p and q such that

$$\phi_{pq} \neq 0, \quad \lambda_p - \lambda_q \neq 0.$$

First we prepare three lemmas.

Lemma 3. $\alpha\alpha_1 - (\alpha\gamma)_1 = 0$.

Proof. From (5.1) we have

$$(\alpha\lambda_p + c)(h - \lambda_p - \lambda_q) = 0,$$

$$(\alpha\lambda_q + c)(h - \lambda_p - \lambda_q) = 0.$$

Thus if $h - \lambda_p - \lambda_q \neq 0$, then we have $\lambda_p = \lambda_q = 0$, which is a contradiction. Hence we have $\alpha h - \alpha\lambda_p - \alpha\lambda_q = 0$. Taking account of the coefficient of θ_1 in the exterior derivative of this, it follows from (4.16) that

$$(5.2) \quad (\alpha h)_1 = 0.$$

From (4.26) we have $(\alpha \sum_p \lambda_p)_1 = 0$. Combining this equation with $h = \alpha + \gamma + \delta + \sum_p \lambda_p$, we have

$$(\alpha(h - \alpha - \gamma - \delta))_1 = 0.$$

Eliminating h from this Lemma 3 follows from (5.2). $\square$

Lemma 4. $\alpha_1 \neq 0$ and $\alpha + \gamma - \delta = 0$.

Proof. From (4.24) we have $2\beta\beta_1 - (\alpha\gamma)_1 = 0$. Hence it follows from Lemma 4 that

$$(5.3) \quad \alpha\alpha_1 + \beta\beta_1 = 0.$$

On the other hand, by (4.32) and (4.34) we have $(\gamma - \delta)\delta_1 - \beta\delta_2 = 0$, and therefore $(\gamma - \delta)\alpha_1 - \beta\alpha_2 = 0$. Thus we have

$$(5.4) \quad (\alpha + \gamma - \delta)\alpha_1 = 0 \quad \text{and} \quad (\alpha + \gamma - \delta)\beta_1 = 0.$$

Suppose that

$$(5.5) \quad \alpha_1 = 0.$$

It follows from (3.8) that

$$(5.6) \quad (\beta^2 - \alpha\gamma - c)_3 = 2\beta\beta_3 - \gamma\alpha_3 - \alpha\gamma_3 = 0.$$

From (4.19)–(4.21), (4.23) and (3.8) we have the following:

$$(5.7) \quad \delta\gamma + \beta X_2 - (\gamma - \delta)X_1 = 0,$$

$$(5.8) \quad \alpha_3 + 3\beta\delta - \alpha\beta + \beta X_1 = 0,$$

$$(5.9) \quad \beta_3 + (\gamma - \delta)X_1 + \gamma\delta - \alpha\gamma - c = 0,$$

$$(5.10) \quad \gamma_3 - 2\beta\delta + (\gamma - \delta)X_2 + \beta\gamma = 0.$$

Substituting of (5.8)–(5.10) into (5.6) we have

$$(5.11) \quad X_1 = 4\alpha,$$

by virtue of (3.11) and (5.7). Substituting of this equation into (5.7)–(5.9) we have

$$(5.12) \quad \beta X_2 = 4\alpha(\gamma - \delta) - \delta\gamma,$$

$$(5.13) \quad \alpha_3 + 3\beta\delta + 3\alpha\beta = 0,$$

$$(5.14) \quad \beta_3 + 3\alpha\gamma - 3\alpha\delta + \gamma\delta = 0.$$

It follows from (4.23), (3.8) and (5.12) that

$$(5.15) \quad \alpha\gamma_3 + \beta(3\alpha\gamma - 6\alpha\delta - \gamma\delta) = 0.$$

From (4.22), (5.3) and (5.5) we have $X_3 = 0$ and $\beta_1 = 0$ and therefore $\alpha_2 = \delta_2 = 0$ because of (4.17). Hence, by (3.8) and (5.5), we have $\gamma_1 = 0$, and so $\beta_2 = 0$.

Now put $F = \alpha$ and β in Lemma 1. Then we have

$$\alpha_3(\gamma + X_1) = 0, \quad \beta_3(\gamma + X_1) = 0.$$

If $\gamma + X_1 \neq 0$, then we have $\alpha_3 = \beta_3 = 0$. It follows from (4.15) and (4.25) that α, β and δ are constants and that $\alpha_i = \beta_i = 0$ for $i = 1, 2$. Furthermore, by (3.8) we see that γ is a constant. Thus from (5.13)–(5.15) we have

$$\alpha + \delta = 0,$$

$$3\alpha\gamma - 3\alpha\delta + \gamma\delta = 0,$$

$$3\alpha\gamma - 6\alpha\delta - \gamma\delta = 0.$$

Hence, by (3.2) and (3.8) we have $\alpha^2 - c = 0$ and $2\beta^2 + c = 0$, which is a contradiction. Therefore $X_1 = -\gamma$, which, together with (5.11), implies $\gamma = -X_1 = -4\alpha$. Thus it follows from (5.13) that $\gamma_3 = -4\alpha_3 = 12\beta(\delta + \alpha)$. Therefore from (5.15) we have a contradiction $\alpha\delta = 0$. $\square$

Lemma 5. $X_1 = 0$.

Proof. Suppose that $X_1 \neq 0$. Then from (4.29) we have

$$(-\lambda_p\lambda_q + 2c)(\lambda_p + \lambda_q) + 2(\alpha + \gamma)\lambda_p\lambda_q - 2c\gamma = 0.$$

Multiply above equation by α^3 and take account of the coefficient of θ_1 in the exterior derivative of this equation. Then, from Lemma 3 and (4.16) we have

$$\alpha\lambda_p + \alpha\lambda_q - \alpha\gamma + \alpha^2 = 0.$$

Again, taking account of the coefficient of θ_1 in the exterior derivative of above equation, we have $\alpha\alpha_1 = 0$, which is a contradiction.

Consequently, for all p, q such that $\phi_{pq} \neq 0$, we have $\lambda_p = \lambda_q$. We take p, q such that $\phi_{pq} \neq 0$. Then by (4.29) we have

$$(5.16) \quad \beta^2 \lambda_p - (\lambda_p - \gamma)((\lambda_p)^2 - \alpha \lambda_p - c) = 0.$$

Furthermore, from $(q3p)$, (4.28) and (4.16) we have

$$(\lambda_p)_1(\lambda_p - \delta) = 0.$$

It follows from (4.16) and Lemma 4 that $(\lambda_p)_1 \neq 0$. Hence we have $\lambda_p = \delta$, which, together with (3.8), (3.11) and (5.16), implies $\alpha + \delta = 0$. Therefore α and δ are constants, which is a contradiction. $\square$

Taking account of the coefficient of θ_3 in the exterior derivative of equation $\alpha + \gamma - \delta = 0$ and making use of (4.19), (4.20), (4.23) and Lemma 5, we have

$$(5.17) \quad \alpha(\gamma - \delta)(\gamma(\alpha - \delta)) + \beta^2(\alpha^2 + \alpha\delta - 3\delta^2) = 0.$$

Again taking account of the coefficient of θ_1 in the exterior derivative of (5.17) and taking account of (3.2) and (5.3), we have

$$(5.18) \quad \alpha^4 + 4\alpha^2 c - 2\delta^2 c + 3c^2 + \beta^2(\alpha^2 + 3\delta^2) = 0.$$

Eliminating β from (5.17) and (5.18), we have a polynomial $4\alpha^4 + 7c\alpha^2 + 4c = 0$ of degree four with respect to α . This shows that α is a constant, which contradicts Lemma 4. Consequently we proved $\beta = 0$.

Since (2.6) and $\beta = 0$, we see that α is a constant in M (see [6]). Thus from (3.1) our assumption $\Xi_{ij;1} = 0$ is equivalent to $\alpha h_{ij;1} = 0$. Put $j = 1$ in (2.3). Then by above equation we have $\alpha h_{i1;k} = -c\alpha\phi_{ik}$. Therefore since (2.1) and $d\xi_i = 0$, we have

$$\alpha \sum_{k,l} h_{ik} \phi_{lk} h_{kj} + \alpha^2 \sum_k \phi_{ki} h_{kj} = -\alpha h_{i1;j} = c\alpha\phi_{ij},$$

which implies that $\alpha^2(\phi H - H\phi) = 0$. Owing to Okumura's work or Montiel and Romero's work stated in Introduction, we complete the proof of our Main Theorem.

Remark 3. In previous papers [4] and [5], we classified real hypersurfaces which satisfy $\nabla_\xi R_\xi = 0$ and some several conditions in $M_n(c)$, $c \neq 0$, $n \geq 3$. Combining these results we have the following:

Let M be a real hypersurface in a complex space form $M_n(c)$, $c \neq 0$, $n \geq 3$ which satisfies $\nabla_\xi R_\xi = 0$. Then the following conditions are mutually equivalent.

- (i) M is locally congruent to type A.
- (ii) $R_\xi S = S R_\xi$.
- (iii) $R_\xi \phi S = S \phi R_\xi$.
- (iv) $R_\xi \phi S = R_\xi S \phi$.

Remark 4. Recently the second author extended Main Theorem to the case $n = 2$ [8].

References

- [1] J. Berndt, *Real hypersurfaces with constant principal curvatures in complex hyperbolic spaces*, J. Reine Angew. Math. **395** (1989), 132–141.
- [2] T. E. Cecil and P. J. Ryan, *Focal sets and real hypersurfaces in complex projective space*, Trans. Amer. Math. Soc. **269** (1982), no. 2, 481–499.
- [3] U.-H. Ki and H. Kurihara, *Real hypersurfaces with cyclic-parallel structure Jacobi operators in a nonflat complex space form*, Bull. Aust. Math. Soc. **81** (2010), no. 2, 260–273.
- [4] U.-H. Ki, H. Kurihara, S. Nagai and R. Takagi, *Characterizations of real hypersurfaces of type A in a complex space form in terms of the structure Jacobi operator*, Toyama Math. J. **32** (2009), 5–23.
- [5] U.-H. Ki, H. Kurihara, and R. Takagi, *Jacobi operators along the structure flow on real hypersurfaces in a nonflat complex space form*, Tsukuba J. Math. **33** (2009), no. 1, 39–56.
- [6] U.-H. Ki and Y. J. Suh, *On real hypersurfaces of a complex space form*, Math. J. Okayama Univ. **32** (1990), 207–221.
- [7] M. Kimura, *Real hypersurfaces and complex submanifolds in complex projective space*, Trans. Amer. Math. Soc. **296** (1986), no. 1, 137–149.
- [8] H. Kurihara, *The structure Jacobi operator for real hypersurfaces in the complex projective plane and the complex hyperbolic plane*, Tsukuba J. Math. **35** (2011), 53–66.
- [9] M. Montiel and A. Romero, *On some real hypersurfaces of a complex hyperbolic space*, Geom. Dedicata **20** (1986), no. 2, 245–261.
- [10] M. Okumura, *On some real hypersurfaces of a complex projective space*, Trans. Amer. Math. Soc. **212** (1975), 355–364.
- [11] M. Ortega, J. D. Pérez, and F. G. Santos, *Non-existence of real hypersurfaces with parallel structure Jacobi operator in nonflat complex space forms*, Rocky Mountain J. Math. **36** (2006), no. 5, 1603–1613.
- [12] J. D. Pérez, F. G. Santos, and Y. J. Suh, *Real hypersurfaces in complex projective spaces whose structure Jacobi operator is D-parallel*, Bull. Belg. Math. Soc. Simon Stevin **13** (2006), no. 3, 459–469.
- [13] R. Takagi, *On homogeneous real hypersurfaces in a complex projective space*, Osaka J. Math. **10** (1973), 495–506.
- [14] ———, *Real hypersurfaces in a complex projective space with constant principal curvatures I, II*, J. Math. Soc. Japan **27** (1975), 43–53, 507–516.

U-HANG KI
 DEPARTMENT OF MATHEMATICS
 KYUNGPOOK NATIONAL UNIVERSITY
 DAEGU 702-701, KOREA
E-mail address: uhangki2005@yahoo.co.kr

HIROYUKI KURIHARA
 DEPARTMENT OF LIBERAL ARTS AND ENGINEERING SCIENCES
 HACHINOHE NATIONAL COLLEGE OF TECHNOLOGY
 HACHINOHE, AOMORI 039-1192, JAPAN
E-mail address: kurihara-g@hachinohe-ct.ac.jp