Cloning and Expression in *Pichia pastoris* of a New Cytochrome P450 Gene from a Dandruff-causing *Malassezia globosa*

Eun Chang Lee¹, Seul Ong Ohk¹, Bo Young Suh¹, Nahee Park¹, Beom Joon Kim²,
Donghak Kim³ and Young-Jin Chun¹

¹College of Pharmacy,

²Department of Dermatology, College of Medicine, Chung-Ang University, Seoul 156-756

³Department of Biological Sciences, Konkuk University, Seoul 143-701, Korea

(Received February 19, 2010; Revised February 24, 2010; Accepted February 26, 2010)

The *Malassezia* fungi are responsible for various human skin disorders including dandruff and seborrheic dermatitis. Of the *Malassezia* fungi, *Malassezia globosa* (*M. globosa*) is one of the most common in human scalp. The completed genome sequence of *M. globosa* contains four putative cytochrome P450 genes. To determine the roles of *Malassezia* P450 enzymes in the biosynthesis of ergosterol, we isolated MGL3996 gene from *M. globosa* chromosomal DNA by PCR. The MGL3996 gene encodes an enzyme of 616 amino acids, which shows strong similarity with known CYP52s of other species. MGL3996 gene was cloned and expressed in *Pichia pastoris* (*P. pastoris*) heterologous yeast expression system. Using the yeast microsomes expressing MGL3996 protein, a typical P450 CO-difference spectrum was shown with absorption maximum at 448 nm. SDS-PAGE analysis revealed a protein band of apparent molecular weight 69 kDa and Western blot with anti-histidine tag antibody showed that MGL3996 was successfully expressed in *P. pastoris*. Cloning and expression of a new P450 gene is an important step to study the P450 monooxygenase system of *M. globosa* and to understand the role of P450 enzymes in pathophysiology of dandruff.

Key words: CYP52, Dandruff, *Malassezia* fungi, P450, *Pichia pastoris*

INTRODUCTION

Malassezia species has been known as major pathogenic yeasts that associate with the common skin disorders including dandruff, pityriasis versicolor, seborrheic dermatitis, psoriasis, and atopic dermatitis in human (Guillot *et al.*, 2008; Zisova, 2009; Cafarchia and Otranto, 2008). So far, the genus *Malassezia* has been shown to comprise at least 13 species based on ribosomal DNA characterization and their ability to grow in certain media: *M. furfur*, *M. sympodialis*, *M. slooffiae*, *M. obtusa*, *M. globosa*, *M. restricta*, *M. pachydermatis*, *M. yamatoensis*, *M. nana*, *M. japonica*, *M. equine*, *M. caprae*, and *M. dermatis* (Guého *et al.*, 1996; Mayser *et al.*, 1997; Morishita *et al.*, 2006; Sugita *et al.*, 2005).

Of these *Malassezia* species, *M. globosa* and *M. restricta* are the most commonly isolated species from the skin of dandruff patients (Gemmer *et al.*, 2002). Despite their asso-

ciation with multiple skin disorders, little is known about these yeasts. Recently, total genomic studies of *Malassezia globosa* CBS 7966 were completed and the 8.9 Mb genome, secretory proteome, and expression of selected genes were described (Xu *et al.*, 2007).

Cytochrome P450 (P450, CYP) enzymes are a superfamily of heme-containing monooxygenases involved in the oxidative metabolism of wide range of endogenous and xenobiotic chemicals. P450s also play important roles in the biosynthesis of antibiotics and other biologically active molecules in bacteria, fungi and plants as well as in animals (Nelson *et al.*, 1996). Although the information about P450 enzymes in *Malassezia* species are completely unknown, the role of P450 in the biosynthetic pathway of fungal steroid may be quite crucial for survival of *Malassezia* fungi because azole antifungal agents such as ketoconazole are often used for treating dandruff. The main target enzyme of azole antifungal agent is known as CYP51, sterol 14 α -demethylase in fungi. Exposure of fungi to azole antifungal agents causes depletion of ergosterol and accumulation of 14 α -methylated sterols (Koltin and Hitchcock, 1997; Sheehan *et al.*, 1999). This activity of azoles interferes with the func-

Correspondence to: Young-Jin Chun, College of Pharmacy, Chung-Ang University, 221 Huksuk-dong, Dongjak-gu, Seoul 156-756, Korea
E-mail: yjchun@cau.ac.kr

tions of ergosterol in fungal membranes and disrupts the structure of the membrane and other functions (Georgopadakou and Walsh, 1996) and causes inhibition of fungal growth. Thus, the studies on *Malassezia* P450s are required to understand *Malassezia* biology and to develop specialized anti-dandruff agents.

The genome project of the *M. globosa* CBS 7966 has provided valuable genetic information about *Malassezia* P450s. Protein sequence similarity analysis using BLAST showed that four probable cytochrome P450 genes exist in *M. globosa*. However, their endogenous roles and contribution to *Malassezia* biology remain completely unknown. In the present studies, we describe the cloning and expression in *Pichia pastoris* of the *M. globosa* MGL3996 gene, a new P450 gene encoding a putative protein highly homologous to the CYP52, which is known as n-alkane and fatty acid hydroxylase (Ohkuma *et al.*, 1995; Scheller *et al.*, 1996).

MATERIALS AND METHODS

Materials. Restriction enzymes and T4 DNA ligase were purchased from TaKaRa Bio (Otsu, Shiga, Japan). Zeocin and yeast nitrogen base was purchased from Invitrogen (Carlsbad, CA, USA). Antibody against His-tag was purchased from Cell Signaling Technology (Danvers, MA, USA). HRP-conjugated mouse secondary antibody was from Santa Cruz Biotechnology (Santa Cruz, CA, USA). Other chemicals were of the highest grade commercially available.

Cell culture. *M. globosa* strain was purchased from the American Type Culture Collection (ATCC, Manassas, VA). For broth culture, inocula were prepared by the culture of *M. globosa* on modified Dixon's agar for five to seven days. Then, the colonies were transferred on modified Leeming-Notman medium containing 0.1% glucose, 0.1% peptone, 0.8% bile salts, 0.2% yeast extract, 0.1% glycerol, 0.5% Tween 60, 3% olive oil, and 50 µg/ml chloramphenicol (Riciputo *et al.*, 1996). The growth of yeast at 32°C was monitored at 530 nm, every 24 h for five days.

Construction of P450 expression plasmid. *M. globosa* genomic DNA was extracted using the QIAamp DNA Mini kit (Qiagen). The gene encoding MGL3996 was amplified by PCR from *M. globosa* genomic DNA by using two gene-specific primers (forward primer MGL3996: 5'-CGGTTCGAAATGGCAGCATCCGCGTTCATC-3', initial codon underlined, *Sfi*I site italicized; reverse primer MGL3996: 5'-CGGTCTAGAAAGCTTCTAGTGATGGTGATGGTG-ATGTTAGTTCAATGGCGTCAT-3', stop codon underlined, *Xba*I site italicized, nucleotide sequence encoding six histidine residue bolded). PCR reaction was carried out with genomic DNA as a template, 0.2 mM dNTPs, 1 pmole of each primer, 1.25 mM MgCl₂, 10 × polymerase buffer, and 2.5 U of *Taq* DNA polymerase in a total volume of 50 µl.

The reaction mixture was preheated to 95°C for 10 min and then subjected to 25 cycles of 95°C for 1 min, 62°C for 1 min, and 72°C for 2 min; followed by one final cycle of 72°C for 10 min. PCR products were analyzed by 1% agarose gel electrophoresis. PCR products were double digested with *Sfi*I and *Xba*I, purified, and then, ligated into the restriction sites of the *Pichia pastoris* expression vector pPICZαA (Invitrogen). *E. coli* DH5α was transformed with the expression vector and the positive clones were selected on LB agar plates containing ampicillin. Recombinant plasmid was isolated and analyzed by DNA sequencing.

Expression of P450 in *P. pastoris*. YPD growth medium containing yeast extract (1%), peptone (2%) and glucose (2%) was inoculated with single colonies of *P. pastoris* (200 ml). The cells were cultured at 29°C for 36 h. The cells were then collected by centrifugation for 10 min at 3000 ×g, 4°C and resuspended to an approximate OD₆₀₀ of 2 in 300 ml of BMM medium (1% yeast extract, 2% peptone, 100 mM potassium phosphate, pH 6.0, 1.34% yeast nitrogen base, and 400 ng/ml biotin) supplemented with 0.5 µM δ-aminolevulinic acid, trace element (250 µl/liter) and zeocin (100 ng/ml). Cells were grown at 29°C while shaking at 200 rpm and induced for 96 h by adding methanol (0.5%) every 24 h. After 96 h, the cells were harvested by centrifugation at 3000 ×g for 10 min 4°C. Cells were resuspended in breaking buffer (50 mM sodium phosphate, pH 7.4, 1 mM EDTA, 5% (v/v) glycerol, 2 mM DTT, and 1 mM protease inhibitor). The cell suspension was mixed with an equal volume of acid washed glass beads (0.5–0.75 mm in diameter) and disrupted by vortexing (830 s at 4°C with cooling on ice for 30 s between the cycles). The lysate was separated from cell debris and glass beads by centrifugation at 12,000 ×g for 10 min at 4°C. The supernatant was centrifuged at 100,000 ×g at 4°C for 1 h and then the microsomal pellet was resuspended in breaking buffer and stored at –80°C.

Quantification of cytochrome P450 content. Protein concentrations were estimated using the bicinchoninic acid method according to the supplier's recommendations (Pierce Chemical Co., Rockford, IL, USA) using bovine serum albumin as a standard. Cytochrome P450 concentration in the isolated membranes was determined by the spectral method of Omura and Sato (1964). The membranes were diluted with sodium phosphate buffer (0.1 M, pH 7.4) containing glycerol (10%) and triton X-100 (0.5%, v/v). After adding small pieces of sodium dithionite crystals, a reference spectrum was recorded from 400 to 500 nm using UV-1650PC UV/Visible spectrophotometer (Shimadzu, Japan). The solution was then saturated with carbon monoxide for 60 s and the spectrum was measured again. The cytochrome P450 concentration was calculated from the difference in absorbance between 450 nm and 500 nm with an extinction

SDS-PAGE and western blot. Extracted proteins were separated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) on 10% polyacrylamide gels. Electrophoresis was performed at 40 mA and 100 V. After finishing electrophoresis, SDS-PAGE gel was stained with Coomassie Brilliant Blue solution (0.1% w/v). For Western blot, gel was electrophoretically transferred onto polyvinylidene difluoride (PVDF) membrane. Membranes were blocked in 5% (w/v) nonfat dried milk in Tris-buffered saline with 0.1% tween 20 (TBST) for 1 h at 4°C. Membranes were then incubated for overnight with anti-his-tag antibody at a 1 : 1000 dilution in TBST. Membrane was incubated with a horseradish peroxidase (HRP)-conjugated goat anti-mouse IgG antibody. Proteins were visualized by an enhanced chemiluminescence method using ChemiDoc XRS (Bio-Rad, CA, USA).

RESULTS AND DISCUSSION

Malassezia fungi have been considered as a major cause of dandruff (Faergemann, 1997). Azole antifungal agents such as ketoconazole, itraconazole, fluconazole, or voriconazole have been leading agents used to treat fungal infections of plants, animals, and human. Azole antifungal agents are also valuable for treating dandruff and seborrheic dermatitis. Because P450 enzymes play an important role in fungal ergosterol biosynthesis and CYP51 is considered as a major drug target for azole antifungal agent, we were interested in P450s in *Malassezia* species including *M. globosa*.

Protein sequence similarity analysis using BLAST showed that four probable P450 genes are existed in *M. globosa* genome. Identification of probable P450 genes within a genome is facilitated by the presence of consensus sequences within the amino acid sequences that are known to be present in all P450 proteins. Amino acid sequences such as EXXR in the K-helix and the cysteine residue which forms a fifth axial ligand to the heme iron are important for P450 function and conserved throughout the P450 superfamily (Table 1). A glycine residue, four amino acids upstream of the conserved cysteine in heme binding motif and a threo-

Table 1. Consensus P450 amino acid sequences in *Malassezia globosa*. Residue conserved in the I-helix (Thr), the K-helix (Glu and Arg), and the heme-binding site (Gly and Cys) are highlighted in bold

Gene name	I-helix	K-helix	Heme binding motif
MGL2415	³⁰³ AGQHT ³⁰⁷	³⁶⁵ ETLR ³⁶⁸	⁴⁷¹ FGAGRHCIG ⁴⁸⁰
MGL3996	³⁸³ AGHET ³⁸⁷	⁴⁵⁵ ESVR ⁴⁵⁸	⁵⁴⁸ FLSGPRSCVG ⁵⁵⁷
MGL0310	³³² ASQDA ³³⁶	³⁹¹ ESLR ³⁹⁴	⁴⁶⁷ FGSGPHNCIG ⁴⁷⁶
MGL1059	²⁶¹ AGHET ²⁶⁵	³³³ ECLR ³³⁶	⁴¹⁵ FLMGPRACIG ⁴²⁴

nine residue in the I-helix known to be involved in oxygen activation were also highly conserved. All of four probable P450s have these conserved amino acid sequences characteristics of P450 enzyme.

Nucleotide and amino acid sequence of the MGL3996 were obtained from NCBI database (XM_001728777) and also were confirmed by DNA sequencing of cloned gene from *M. globosa* genomic DNA by PCR (Fig. 1). The MGL3996 gene has 1851 nucleotides and encodes a protein of 616 amino acids which is highly homologous to the CYP52s from other yeast species. Amino acid sequence alignment of MGL3996 with *Candida albicans* CYP52A1, *Candida tropicalis* CYP52A8, *Pichia stipitis* CYP52A3 and

1 atggcagcattcgggttcattctcgggtgcagatgtgtctctaacctcgtcgggtctggcg
61 M A A S A S A F I F S G A D V L L Y L L A A G L A
111 ggtctctcagtttgtctgtacccgtcgtttctcatcttgaacagctgttgcaagcgcctctac
161 G L L V A V P L L F L I L L N Q L V Q A L Y
211 cgcattgaccatttgggtgagctgatcaaccccttgggaagcacatacctaaggtgaagaccact
261 R M T I G E L I N P W K H I F K V K T T
311 gatcctctgtccctcatctctcgggtgattttctgtctatcaaaagctgacagccctgctcaa
361 D P L S L I F G A G D F R A I K A A D D P A Q
411 cagcatctcatctggatgaacaaagcttggccaagctgaccgcctaccgcctatgtgtttac
461 Q H V I M N N K L G Q V Y T R R H M F Y
511 gtgcagcgtgcttcttctgtctgtaccaagcgctaatcaacgtgttgagccctgggaaa
561 G C R V L L L A D P K A L I H V L S P A K
611 vtgtaccagtatcccaagccgaatatacagcagcagctttcttcggcagctgtacttgagat
661 A Y Q Y P K P E Y T S S F L A A V L L G D
711 ggtctgtcatctcaggggtgaaaagcagcagccgcaaaaactcattgcgcctgca
761 G L V S I E G E K H A R Q R K I I A P A
811 ttgtgcgaagctgttctaaaagactaccagccaaactcagcgtcatgtcaagctgctt
861 F A P S V V K D Y Q P T T I H R H A K L L
911 gtgcagacgtcctcgtcatgtgcaacaggaaatlgatcgcacagctggcgaagctcccg
961 V E K L R L I V Q Q A E I D R H S G K I P
1011 gtgcacactcacattactcggcagctcagaatctgacgcagcagctgtgattgaagaacct
1061 A D T H I T G Q S E S E D R S V V L K N P
1111 ggaagctgaagctgtgatlgatgacatctctggtctcttcgtactcagctcgcagatcatl
1161 G S E A V I D V L F W L S R T T L D I I
1211 ggtgaaattgggtttgatgcgaacttcggctgccttgagcgtggaacccgtgaccgcgtt
1261 G E I G F D A D F R S L E R G N R D P L
1311 gcgaatlcattcggcgagcgcacagggagcagctccllaacatlgacncttgcgttgcgtc
1361 A N S M A T L M E A V A L N I D L G L A I
1411 ttcttatctcttctgaaagccgcgttctcagctgcctcgtgaagatctcgacgaagcgt
1461 F L I L S E K P G L Q W L R K I P T K R
1511 actacgcaactcgtgtatcgaagaagcgttgcgaagcagctgcagaagaactgtgcag
1561 T T T I R D A K T K T A V E K H A E I V E
1611 cgcactgcgaatgagatctcttctgaaagctcaagctctgtaacgaagaaacttgacgat
1661 R M R N E I L S E S Q G L S K E N F D D
1711 ggtcgtcgtcagggtgcgaagtccatacttcggctgtatgacctgcacaaactggtctcg
1761 G S A T R S K S L I S R M I R A N M A S
1811 gacctgaacatcacaagcgcatctcgaatcgtgaattaatggctcagatgacqactttg
1861 D L K P S E R M S N A E L M G Q M T T L
1911 atcatcgttcggccagcagaactgcacactcgaagcagctggggccctttgctttggcc
1961 I I A G H E T T A T Q N S W L L W L L A
2011 atgcaacctgagggcagaagactctctcgttaaggagctccagaagctgtggcgcaagaa
2061 M H P E A Q D R L R L K E A V A K E
2111 cggcgcgaaatggacaactctgtcggaggagggaactgccatctatgacacccagccgtt
2161 R A E M D N L S E E E R A I Y D T Q P V
2211 cqqatgatacaactcgttaccatctctcgacaacgcttgatgaagaagagatgtcgatcatgctt
2261 R D I N S L P Y L D N V K E S V R M L
2311 cccctgccttcctgacactgtgcgtctgcctcgaagtgatgatgctccctctgtcctgt
2361 P S I P S T V R V A L K D D V P L S R
2411 gqgttacaagcgcgctgatgtatgcgtatcagctacaactcgatcatgatccaaagdgctac
2461 G Y K R A D G K G T Y N S I M I P K G H
2511 gagctctttatccccctgaatvgatccaactctcgaaggaaactatgggggtgaagatgct
2561 E L F I P L A T V I Q L S K E L W G E D A
2611 caaagattcaaccccagcgttgggacaatttggccctctccgtgtacatgaatgcaaaagt
2661 Q D F N P S R W D N I L P S S V T N A K M
2711 ccaocgggtcaccctgtcgtctttctcgaagcaactgcctctcgtgttgccaagcagaca
2761 P F G H L F A F L S G P R S C V G K Q T
2811 gctattctcgaaacacaggtcatctcggccgcatagtatctcaacttccgcttctgacgtt
2861 A I L E T V Q I L A H M I L N F R F D V
2911 gtgcagcgtgggattcgtctcaagcgcagagattgttgcgtcgtgccttcgtcgaaggc
2961 V F G W D L V Q R Q Q I V R A F V E G
3011 caatgggaaggagcactcgttatgccactcattatgacgcacttaactaa
3061 Q W K E G I R M P L I M T P L N -

Fig. 1. Nucleotide sequence of *Malassezia* MGL3996 gene and the predicted amino acid sequence. Nucleotide sequence of MGL3996 were obtained from NCBI database (XM_001728777) and was confirmed by nucleotide sequencing of PCR-cloned gene from *M. globosa* genomic DNA. Nucleotide numbering and amino acid numbering are shown at thin and bold, respectively.

M. globosa	MAASAF-----ISGADVLYLAAGLAGLVAVPLLFILNQVLQVQALYRMTIGELINPWKHIP	M. globosa	AEEIVERMRNEILSESQGL-----SKENFDDGSATRSKSLISRMIRANMASDLKPSERMSN
C. albicans	MSPSLVHQILAITHPYVEYLEENITKWYILVPGILIVNLVTTLL-----	C. albicans	SDFYVQKALSQDELDEQ-----KGVVFLY-----ELAKQTRD-F
C. tropicalis	MYEQV-----VEYVYVPLVPLVILHKVDFDMMH-----	C. tropicalis	TNYVQALDAPTEELEKQ-----SGVVFY-----ELVKQTRG-F
P. stipitis	MF-----	P. stipitis	ARFYVDALNFSQAELEKKS--QESYDTLL-----ELVQTRD-F
L. bicolor	MDVPP-----GIAYLVQRLPRILTPPTLAYALVHL-----GKTHLS	L. bicolor	IDPIVAAASRKKEMLSPSLDRDDETL-----ELVNSTED-F
	*		* *
M. globosa	KVK-TTDSLIFGDF-----R-	M. globosa	AEIMGQMTTLIIAGHETTATQNSWALWLLAMHPEAQDRLLKEELQEAQVAK-ERAEMDNLSE
C. albicans	-----TKYLEYKFN-----KP	C. albicans	KVLRDQLNILLVAGRDITAGLSFVFFELARNPEVYAKLKEEYKNGFSGEDARIDE---
C. tropicalis	-----TRRLMKQLGA-----AP	C. tropicalis	NVLRDQSLNILLVAGRDITAGLSFVFFELARNPEVYAKLKEEYKNGFSGEDARIDE---
P. stipitis	-----	P. stipitis	KVLQDQLAILMAGRDITAGLSFVFFELARNPEVYAKLKEEYKNGFSGEDARIDE---
L. bicolor	IELSTWGLSSVLLSLPGLTSLVWKDLVIRWASNGKAVLPVPAPDFPGGARSLYRA	L. bicolor	IILRDEIMSLVAGRDITAGLSFVFFELARNPEVYAKLKEEYKNGFSGEDARIDE---
			* . : : : * : : : : : : : : : * : : : : . .
M. globosa	--AIKAADP-AQQ--HVIWMKLGQVYR-----YRH-MFYVQRV	M. globosa	EERAIYDTQVPRDINSPLVDNVHESVMLPSIPSTVRVAKDKDVPVLSRGYKRDGKG
C. albicans	VTNFIQDYTFGVITPLILYKSGQVMEFANFNWFKLIKPNVDTGTELR-IFGMHLI	C. albicans	-----ITFESLKQCEYLKAVINESLPLVSPVPHNFRATRTNTTLPRGG-----GPD
C. tropicalis	VTNQLDHNFNGIINGWKHLSSKK--VELKN-IMIINLPKRFQVWVHMLVPSLEQSS	C. tropicalis	-----ITFESLKQCEYLKAVINESLPLVSPVPHNFRATRTNTTLPRGG-----GSD
P. stipitis	-----	P. stipitis	-----ITFESLKQCEYLKAVINESLPLVSPVPHNFRATRTNTTLPRGG-----GED
L. bicolor	VTSFKTGYP-GVST-LILKVDG-----LQDMCKFGGYTF-----MSK-ILFEYRV	L. bicolor	-----PTYDDFKEMKFLKAVINETLPLVSPVPHNFRATRTNTTLPRGG-----VP
			. . : : : . : : * : * : * : * : *
M. globosa	LLADPKALIHVSPAKAYQPKPEYTS-FLAAYLGDGLVSEIEGKHAR-----	M. globosa	TYNSIMIPKGHELFILNVLQSLKELGADA-----QDFNPSRWDN
C. albicans	ETKDPENIKAVLA-TQFNDFSLG-TRHG-FLYSLLGDGIFTLDGAGWKH-----	C. albicans	GMSPVIVKKGQSVMYTVLTHRTKTYTGADA-----NEFFPERWFE
C. tropicalis	SQKIRRNKALLA-TQFNDFSLG-KRHT-LFKPLLDGIFTLDGAGWKH-----	C. tropicalis	GMSPVIVKKGQSVMYTVLTHRTKTYTGADA-----NEFFPERWFE
P. stipitis	MTTDPENFKAMLA-TQFNDFSLG-KRHT-LFKPLLDGIFTLDGAGWKH-----	P. stipitis	GKSPFIPRGTTVGFRVYSTQNKKEYGDEP-----DVFFPERWAD
L. bicolor	VTAEPENIKAMLA-TQFNDFSLG-KRHT-LFKPLLDGIFTLDGAGWKH-----	L. bicolor	GKSPFIPRGTTVGFRVYSTQNKKEYGDEP-----DVFFPERWAD
	. : : * : : : : : : : : * : : : : *		. . : : : . : : * : * : * : * : *
M. globosa	-----QRKIAPAFAPSVVKDYQPTIHRHAKLLVEKLRLVQQEIDRHSKIPAD	M. globosa	LPSSVINAKMPPGHFLFSLGSPSCVSKQTALLETQVILAHMILNFRFVDVPG-WDL---
C. albicans	-----SRMLRPQFAREQIAHVK-ILEPHIQVFFKHI-----	C. albicans	PETRLK-----G-WAYLHFNNGSPRICLQGFALTEAGYVLRVLAQSFDTLELAPFVYFP-
C. tropicalis	-----SRMLRPQFAREQIAHVK-ILEPHIQVFFKHI-----	C. tropicalis	PETRLK-----G-WAYLHFNNGSPRICLQGFALTEAGYVLRVLAQSFDTLELAPFVYFP-
P. stipitis	-----SRMLRPQFAREQIAHVK-ILEPHIQVFFKHI-----	P. stipitis	I--GKL-----G-WAYLHFNNGSPRICLQGFALTEAGYVLRVLAQSFDTLELAPFVYFP-
L. bicolor	IFLTSLRVFRHRTMPFFSKDRISHFD-NFGRHADDALGQLKA-----	L. bicolor	ERLHKYLTNP-NFLEFNAGSPRICLQGFALTEAGYVLRVLAQSFDTLELAPFVYFP-
	* : * * . . : : * . : :		. . * . * . * : * : * : * : . . : : *
M. globosa	THITQGESDRSVVLKNGSEAVIDVLEWLSRTTLDIIEIGFADFRSLERGNRDLAN	M. globosa	-----VQRQIVRAAFVGEQWKGIRIMPLIMTLIN
C. albicans	-----RKNKGKT--FDIQELFFRLTVDSSTEFLEFGSVSLRDETIG-----	C. albicans	-----KLMNSLTLSLLDGANV-----EM--Y
C. tropicalis	-----VKNKGKT--FDIQELFFRLTVDSSTEFLEFGSVSLRDETIG-----	C. tropicalis	-----KRLTNLMSLQDGTIV-----KI--D
P. stipitis	-----RSYKGET--VDLQQLFTKFTLDATTEFLFGSVSLRDETIG-----	P. stipitis	-----KRLTNLMSLQDGTIV-----KI--D
L. bicolor	-----RLREGHA--VDQDLISRFTLDATTEFLFGSVSLRDETIG-----	L. bicolor	-----KRLTNLMSLQDGTIV-----KI--D
	* . . : : * : * : * : * : * : *		RAPAIWATEDRTGRKGREKIPKTHLTMVQDGLV-----TMEAK
			: : : : : : : : : : : : *
M. globosa	SMATLMEA--VLNID-----LGLAIFLILSEKPLQWLRIKPTKRTTTIRDAKKTVEKH		
C. albicans	--MSPSVKNLAGRDEFADAFNYSQTINAYRFLQGFYWLGN-----GTFKFKSIAAVHKF		
C. tropicalis	--SYQDDIDFVGRKDEAFNKAQYLAIRTLVQDFYVLVN-----NQFEDCNKLVHKF		
P. stipitis	--MKTDP-DVYAKDFDTGLFITQKVTSEYGAQGFYWLID-----GKEFFRTIANVHKF		
L. bicolor	STTSTSDSDAHPTFRVHAFOEAMITAFRSFGSHWPLFEF--HDMKKFMVYVHDF		
	* . : : : *		

Fig. 2. Multiple sequence alignment of the *Malassezia* MGL3996 amino acid sequence with four other CYP52 proteins. The amino acid sequences were aligned using the software T-Coffee from the Swiss Institute of Bioinformatics (<http://www.ch.embnet.drg/software/Tcoffee.html>). *M. globosa* MGL3996 sequence was compared with *Candida albicans* CYP52A1, *Candida tropicalis* CYP52A8, *Pichia stipitis* CYP52A3 and *Laccaria bicolor* CYP52-1 amino acid sequences. Boxes indicate EXXR region in K-helix and conserved heme-binding motif. (*) indicate identical residues; (:) indicate semi-conserved residues; (.) designate conserved residues.

Laccaria bicolor CYP52-1 revealed a high sequence similarity with a score 78, 78, 84 and 72%, respectively (Fig. 2). The good sequence alignment of MGL3996 with other CYP52 suggests that they may have similar enzymatic specificities.

To confirm whether MGL3996 gene product shows a property of P450 enzymes, cloned MGL3996 gene was subcloned into the pPICZαA expression vector for expressing in *P. pastoris* yeast using restriction sites of *SfuI* and *XbaI* (Fig. 3). To facilitate expression and further purification of P450, six histidine residues were introduced just before the termination codon. The recombinant pPICZαA plasmid was successfully transformed into *P. pastoris* X-33 strain for high-level expression. After the continuous induction with methanol for 96 h, cells were collected and membrane fractions were isolated. Isolated membrane fractions expressing MGL3996 protein had a typical Fe²⁺-CO versus Fe²⁺ difference spectrum which is characteristic for P450 enzymes (Fig. 4). The absorption maximum was at 448 nm and the expression level was about 2 nmol P450 per ml of membranes.

To verify the expression of MGL3996 protein in *P. pastoris*, SDS-PAGE and Western blot analysis were per-

formed with cellular lysates after methanol induction for 96 h. SDS-PAGE result showed that a protein band of apparent molecular weight of 69 kDa was highly induced in the induced cell carrying pPICZαA/MGL3996. No such band was detected in the control cells (Fig. 5A). Western blot with antibody against histidine tag clearly showed the band corresponding in size to the MGL3996 (Fig. 5B).

In this study, we report expression of *M. globosa* P450 in *P. pastoris* for the first time. Previously, we had been tried to express MGL3996 gene in *E. coli* system but could not detect P450 activity. Thus, we attempted to express MGL3996 gene in *P. pastoris* yeast expression system. The *P. pastoris* yeast system is heterologous protein expression systems to produce high levels of functional proteins (Li et al., 2007; Daly and Hearn, 2005), which include membrane-bound proteins (Cregg et al., 1993; Cereghino and Cregg, 2006). The *P. pastoris* yeast system provides several advantages including stability of expression, easy-to-handling or a correct protein folding (Kolar et al., 2007). *P. pastoris* is considered as a better expression host for eukaryotic gene expression compared to *S. cerevisiae*.

The result that MGL3996 produced a clear CO-differ-

Fig. 3. *Malassezia* MGL3996 expression vector in *P. pastoris*. 5' AOX1: alcohol oxidase 1 promoter region allows methanol-inducible expression in *P. pastoris*, Zeocin: zeocin resistance gene derived from *Streptoalloteichus hindustanus*, AOX1 transcription terminator region: native transcription termination and polyadenylation signal from AOX1 gene permits mRNA processing, MGL3996: gene encoding *M. globosa* putative P450.

Fig. 4. Fe^{2+} -CO versus Fe^{2+} difference spectra of *Malassezia* MGL3996 expressed in *P. pastoris*. *P. pastoris* cells transformed with pPICZ α A/MGL3996 were incubated in BMM media for 96 h after methanol induction. The membrane fractions were isolated and Fe^{2+} -CO versus Fe^{2+} difference spectrum was measured. The calculated P450 level in the microsomal fraction is about 1.6 nmol per ml.

ence spectrum indicates that the MGL3996 may be a real P450 enzyme existed in *M. globosa* and *P. pastoris* system produces a correctly folded P450 protein. Although *P. pas-*

Fig. 5. SDS PAGE and Western blot analysis of expressed MGL3996. *P. pastoris* cells transformed with pPICZ α A/MGL3996 were incubated in BMM media for 96 h after methanol induction. Whole cell extracts were isolated, and then 10% SDS-PAGE and Western blot analysis was performed. (A) SDS-PAGE. Lane 1, protein marker; lane 2, lysate of transformed cells; lane 3: lysate of control cells. Indicated 69 kDa protein band represent the expressed MGL3996 protein. (B) Western blot analysis. Western blot was performed with anti-His-tag antibody. Lane 1, lysate of transformed cells; lane 2: lysate of control cells. Indicated 69 kDa protein band represent the MGL3996 protein.

toris may contain endogenous P450 proteins, these must be expressed in very low levels because endogenous P450s were never detectable by CO-difference spectroscopy.

To date, only seven P450 enzymes have been expressed in *P. pastoris*, i.e., spiny dogfish shark CYP17 (Trant, 1996), plant CYP79D1 (Andersen *et al.*, 2000), fungal PcCYP1f (Matsuzaki and Wariishi, 2005), human CYP2D6 (Dietrich *et al.*, 2005), human CYP17 α (Kolar *et al.*, 2007), Arabidopsis CYP78A9 (Ito and Meyerowitz, 2000) and CYP85A2 (Katsumata *et al.*, 2008). For the first time, here we report successful expression of a *Malassezia* P450 in a *P. pastoris* expression system.

Altogether, our findings indicate that MGL3996, a probable P450 gene selected from *M. globosa* genome using sequence similarity analysis may be a functional P450 enzyme although information about enzymatic properties and chemical reactions still needs to be determined. Our establishment of *P. pastoris* expression system for *Malassezia* P450 gene will be valuable to understand the role of other P450 enzymes in *Malassezia* species.

ACKNOWLEDGEMENT

This study was supported by a grant of the Korea Healthcare Technology R&D Project, Ministry of Health, Welfare and Family Affairs (A080065).

REFERENCES

- Andersen, M.D., Busk, P.K., Svendsen, I. and Møller, B.L. (2000). Cytochromes P-450 from cassava (*Manihot esculenta* Crantz)

- catalyzing the first steps in the biosynthesis of the cyanogenic glucosides linamarin and lotaustralin. Cloning, functional expression in *Pichia pastoris*, and substrate specificity of the isolated recombinant enzymes. *J. Biol. Chem.*, **275**, 1966-1975.
- Cafarchia, C. and Otranto, D. (2008). The pathogenesis of *Malassezia* yeasts. *Parassitologia*, **50**, 65-67.
- Cereghino, J.L. and Cregg, J.M. (2006). Heterologous protein expression in the methylotrophic yeast *Pichia pastoris*. *FEMS Microbiol. Rev.*, **24**, 45-66.
- Cregg, J.M., Vedvick, T.S. and Raschke, W.C. (1993). Recent advances in the expression of foreign genes in *Pichia pastoris*. *Bio/Technology*, **11**, 905-910.
- Daly, R. and Hearn, M.T. (2005). Expression of heterologous proteins in *Pichia pastoris*: a useful experimental tool in protein engineering and production. *J. Mol. Recognit.*, **18**, 119-138.
- Dietrich, M., Grundmann, L., Kurr, K., Valinotto, L., Saussele, T., Schmid, R.D. and Lange, S. (2005). Recombinant production of human microsomal cytochrome P450 2D6 in the methylotrophic yeast *Pichia pastoris*. *ChemBiochem*, **5**, 2014-2022.
- Faergemann, J. (1997). *Pityrosporum* yeasts-what's new? *Mycoses*, **40**, 29-32.
- Gemmer, C.M., DeAngelis, Y.M., Theelen, B., Boekhout, T. and Dawson, T.L. Jr. (2002). Fast, noninvasive method for molecular detection and differentiation of *Malassezia* yeast species on human skin and application of the method to dandruff microbiology. *J. Clin. Microbiol.*, **40**, 3350-3357.
- Georgopapadakou, N.H. and Walsh, T.J. (1996). Antifungal agents: chemotherapeutic targets and immunologic strategies. *Antimicrob. Agents Chemother.*, **40**, 279-291.
- Guého, E., Midgley, G. and Guillot, J. (1996). The genus *Malassezia* with description of four new species. *Antonie Van Leeuwenhoek*, **69**, 337-355.
- Guillot, J., Hadina, S. and Guého, E. (2008). The genus *Malassezia*: old facts and new concepts. *Parassitologia*, **50**, 77-79.
- Ito, T. and Meyerowitz, E.M. (2000). Overexpression of a gene encoding a cytochrome P450, CYP78A9, induces large and seedless fruit in arabidopsis. *Plant Cell*, **12**, 1541-1550.
- Katsumata, T., Hasegawa, A., Fujiwara, T., Komatsu, T., Notomi, M., Abe, H., Natsume, M. and Kawaide, H. (2008). Arabidopsis CYP85A2 catalyzes lactonization reactions in the biosynthesis of 2-deoxy-7-oxalactone brassinosteroids. *Biosci. Biotechnol. Biochem.*, **72**, 2110-2117.
- Kolar, N.W., Swart, A.C., Mason, J.I. and Swart, P. (2007). Functional expression and characterisation of human cytochrome P45017A in *Pichia pastoris*. *J. Biotechnol.*, **129**, 635-644.
- Koltin, Y. and Hitchcock, C.A. (1997). Progress in the search for new triazole antifungal agents. *Curr. Opin. Chem. Biol.*, **1**, 176-182.
- Li, P., Anumanthan, A., Gao, X.G., Ilangovan, K., Suzara, V.V., Düzgüneş, N. and Renugopalakrishnan, V. (2007). Expression of recombinant proteins in *Pichia pastoris*. *Appl. Biochem. Biotechnol.*, **142**, 105-124.
- Matsuzaki, F. and Wariishi, H. (2005). Molecular characterization of cytochrome P450 catalyzing hydroxylation of benzoates from the white-rot fungus *Phanerochaete chrysosporium*. *Biochem. Biophys. Res. Commun.*, **334**, 1184-1190.
- Mayser, P., Haze, P., Papavassilis, C., Pickel, M., Gruender, K. and Guého, E. (1997). Differentiation of *Malassezia* species: selectivity of cremophor EL, castor oil and ricinoleic acid for *M. furfur*. *Br. J. Dermatol.*, **137**, 208-213.
- Morishita, N., Sei, Y. and Sugita, T. (2006). Molecular analysis of *Malassezia* microflora from patients with pityriasis versicolor. *Mycopathologia*, **161**, 61-65.
- Nelson, D.R., Koymans, L., Kamataki, T., Stegeman, J.J., Feyereisen, R., Waxman, D.J., Waterman, M.R., Gotoh, O., Coon, M.J., Estabrook, R.W., Gunsalus, I.C. and Nebert, D.W. (1996). P450 superfamily: update on new sequences, gene mapping, accession numbers and nomenclature. *Pharmacogenetics*, **6**, 1-42.
- Ohkuma, M., Muraoka, S., Tanimoto, T., Fujii, M., Ohta, A. and Takagi, M. (1995). CYP52 (cytochrome P450alk) multigene family in *Candida maltosa*: identification and characterization of eight members. *DNA Cell Biol.*, **14**, 163-173.
- Omura, T. and Sato, R. (1964). The carbon-monoxide binding pigment of liver microsomes. *J. Biol. Chem.*, **239**, 2370-2378.
- Riciputo, R.M., Oliveri, S., Micali, G. and Sapuppo, A. (1996). Phospholipase activity in *Malassezia furfur* pathogenic strains. *Mycoses*, **39**, 233-235.
- Scheller, U., Zimmer, T., Kärger, E. and Schunck, W.H. (1996). Characterization of the n-alkane and fatty acid hydroxylating cytochrome P450 forms 52A3 and 52A4. *Arch. Biochem. Biophys.*, **328**, 245-254.
- Sheehan, D.J., Hitchcock, C.A. and Sibley, C.M. (1999). Current and emerging azole antifungal agents. *Clin. Microbiol. Rev.*, **12**, 40-79.
- Sugita, T., Tajima, M., Ito, T., Saito, M., Tsuboi, R. and Nishikawa, A. (2005). Antifungal activities of tacrolimus and azole agents against the eleven currently accepted *Malassezia* species. *J. Clin. Microbiol.*, **43**, 2824-2829.
- Trant, J.M. (1996). Functional expression of recombinant spiny dogfish shark (*Squalus acanthias*) cytochrome P450c17 (17 alpha-hydroxylase/C17,20-lyase) in yeast (*Pichia pastoris*). *Arch. Biochem. Biophys.*, **326**, 8-14.
- Xu, J., Saunders, C.W., Hu, P., Grant, R.A., Boekhout, T., Kuramae, E.E., Kronstad, J.W., Deangelis, Y.M., Reeder, N.L., Johnstone, K.R., Leland, M., Fieno, A.M., Begley, W.M., Sun, Y., Lacey, M.P., Chaudhary, T., Keough, T., Chu, L., Sears, R., Yuan, B. and Dawson, T.L. Jr. (2007). Dandruff-associated *Malassezia* genomes reveal convergent and divergent virulence traits shared with plant and human fungal pathogens. *Proc. Natl. Acad. Sci. USA*, **104**, 18730-18735.
- Zisova, L.G. (2009). *Malassezia* species and seborrheic dermatitis. *Folia Med.*, **51**, 23-33.